

ISSN EXERCISE & SPORT NUTRITION REVIEW: RESEARCH & RECOMMENDATIONS

Richard B. Kreider¹, Anthony L. Almada², Jose Antonio³, Craig Broeder⁴, Conrad Earnest⁵, Mike Greenwood¹, Thomas Incledon⁶, Douglas S. Kalman⁷, Susan M. Kleiner⁸, Brian Leutholtz¹, Lonnie M. Lowery⁹, Ron Mendel¹⁰, Jeffrey R. Stout¹¹, Darryn S. Willoughby¹, Tim N. Ziegenfuss¹⁰

Exercise & Sport Nutrition Lab, Baylor University, Waco, TX¹; IMAGINutrition, Laguna Nigel, CA²; Juvalution, Fort Lauderdale, FL³; Department of Biological Sciences, Clinical Exercise Physiology Program, Benedictine University, Lisle, IL⁴; The Cooper Institute, Dallas, TX⁵; Human Performance Specialists, Inc., Chandler, AZ⁶; Miami Research Associates, Miami, FL⁷; Department of Medical History and Ethics, University of Washington, Seattle, WA⁸; Human Nutrition Laboratory, Department of Nutrition and Dietetics, Kent State University, Kent, OH⁹; Ohio Research Group of Exercise Science & Sports Nutrition, Wadsworth, OH¹⁰; Department of Exercise Science and Health Promotion, Florida Atlantic University, Davie, FL¹¹. Sports Nutrition Review Journal. 1(1):1-44, 2004. Address correspondence to [Richard Kreider@baylor.edu](mailto:Richard.Kreider@baylor.edu).

Received May 10, 2004/Accepted May 15, 2004/Published (online)

ABSTRACT

Sport nutrition is a constantly evolving field with literally thousands of research papers published annually. For this reason, keeping up to date with the literature is often difficult. This paper presents a well-referenced overview of the current state of the science related to how to optimize training through nutrition. More specifically, this article discusses: 1.) how to evaluate the scientific merit of nutritional supplements; 2.) general nutritional strategies to optimize performance and enhance recovery; and, 3.) our current understanding of the available science behind weight gain, weight loss, and performance enhancement supplements. Our hope is that ISSN members find this review useful in their daily practice and consultation with their clients. *Sports Nutrition Review Journal. 1(1):1-44, 2004.*

Key Words: sport nutrition, dietary supplements, ergogenic aids, weight gain, weight loss

INTRODUCTION

Sport nutrition professionals need to know how to evaluate the scientific merit of articles and advertisements about exercise and nutrition products so they can separate marketing hype from scientifically based training and nutritional practices. In order to help educate ISSN members about sport nutrition, we have updated a letter to the Editor ([PEP Online. 6\(10\), 2003](#)) which represents a compilation of Dr. Kreider's published work in this area for the Sport Nutrition Review Journal's inaugural issue. This paper provides an overview of: 1.) what

are ergogenic aids and dietary supplements; 2.) how dietary supplements are legally regulated; 3.) how to evaluate the scientific merit of nutritional supplements; 4.) general nutritional strategies to optimize performance and enhance recovery; and, 5.) an overview of our current understanding of the ergogenic value weight gain, weight loss, and performance enhancement supplements. We have also categorized nutritional supplements into apparently effective, possibly effective, too early to tell, and apparently ineffective as well as describes our general approach to educating athletes about sport nutrition. While some may not agree with all of our

interpretations of the literature and/or categorization of a particular supplement and some classifications may change over time as more research is forthcoming, these interpretations are based on the current available scientific evidence and have been well received within the broader scientific community. Our hope is that ISSN members find this information useful in their daily practice and consultation with their clients.

WHAT IS AN ERGOGENIC AID?

An ergogenic aid is any training technique, mechanical device, nutritional practice, pharmacological method, or psychological technique that can improve exercise performance capacity and/or enhance training adaptations^{1,2}. This includes aids that may help prepare an individual to exercise, improve the efficiency of exercise, and/or enhance recovery from exercise. Ergogenic aids may also allow an individual to tolerate heavy training to a greater degree by helping them recover faster or help them stay healthy during intense training. Although this definition seems rather straightforward, there is considerable debate regarding the ergogenic value of various nutritional supplements. Some sport nutrition specialists only consider a supplement ergogenic if studies show that the supplement significantly enhances exercise performance (e.g., helps you run faster, lift more weight, and/or perform more work during a given exercise task). On the other hand, some feel that if a supplement helps prepare an athlete to perform or enhances recovery from exercise, it has the potential to improve training adaptations and therefore should be considered ergogenic. In our view, one should take a broader view about the ergogenic value of supplements. While we are interested in determining the performance enhancement effects of a supplement on a single bout of exercise, we also realize that one of the goals of training is to help people tolerate training to a greater degree. People who tolerate training better usually experience greater gains from training over time. Consequently, employing

nutritional practices that help prepare people to perform and/or enhance recovery from exercise should also be viewed as ergogenic.

WHAT ARE DIETARY SUPPLEMENTS AND HOW ARE THEY REGULATED?

According to the Food and Drug Administration (FDA), dietary supplements were regulated in the same manner as food prior to 1994³. Consequently, the manufacturing processes, quality, and labeling of supplements were monitored by FDA. However, many people felt that the FDA was too restrictive in regulating dietary supplements. As a result, Congress passed the Dietary Supplement Health and Education Act (DSHEA) in 1994 which placed dietary supplements in a special category of "foods". In October 1994, DSHEA was signed into law by President Clinton. The law defined a "dietary supplement" as a product taken by mouth that contains a "dietary ingredient" intended to supplement the diet. "Dietary ingredients" may include vitamins, minerals, herbs or other botanicals, amino acids, and substances (e.g., enzymes, organ tissues, glandulars, and metabolites). Dietary supplements may also be extracts or concentrates from plants or foods. Dietary supplements are typically sold in the form of tablets, capsules, soft gels, liquids, powders, and bars. Products sold as dietary supplements must be clearly labeled as a dietary supplement.

According to DSHEA, dietary supplements are not drugs. Dietary supplement ingredients that were sold prior to 1994 are therefore not required to be shown to be safe and/or effective in clinical trials prior to being approved for sale by the FDA. However, new dietary supplement ingredients introduced after 1994 must undergo pre-market review for safety data by the FDA before it can be legally sold. Supplement companies are responsible for determining that the dietary supplements it manufactures or distributes are safe and that any representations or claims made about them are substantiated by

adequate evidence to show that they are not false or misleading. Because of this, DSHEA requires supplement manufacturers to include on the label that "This statement has not been evaluated by the FDA. This product is not intended to diagnose, treat, cure, or prevent any disease". According to the 1994 Nutrition Labeling and Education Act (NELA), the FDA has the ability to review and approve health claims for dietary supplements and foods. However, since the law was passed, it has only reviewed a few claims. The delay in reviewing health claims of dietary supplements resulted in a law suit filed by Pearson & Shaw et al v. Shalala et al in 1993. After years of litigation, U.S. Court of Appeals for the District of Columbia Circuit ruled in 1999 that qualified health claims may now be made about dietary supplements with approval by FDA as long as the statements are truthful and based on science. Supplement companies wishing to make health claims about supplements can submit research evidence to the FDA for approval. Additionally, they must submit an Investigation of New Drug (IND) application to FDA if a research study on a nutrient is designed to treat an illness and/or medical affliction and/or the company hopes to one day obtain approval for making a qualified health claim if the outcome of the study supports the claim. Studies investigating structure and function claims, however, do not need to be submitted to the FDA as an IND.

Manufacturers and distributors of dietary supplements are not currently required to record, investigate or forward to FDA any reports they receive on injuries or illnesses that may be related to the use of their products. However, the FDA and other groups have established phone hotlines and online adverse event monitoring systems to report problems they believe may be a result of taking dietary supplements. While these reports are unsubstantiated, can be influenced by media attention to a particular supplement, and do not necessarily show a cause and effect, they are used by the FDA to monitor

trends and "signals" that may suggest a problem. Once a dietary supplement product is marketed, the FDA has the responsibility for showing that a dietary supplement is unsafe before it can take action to restrict the product's use or removal from the marketplace. The Federal Trade Commission (FTC) is responsible to make sure manufacturers are truthful regarding claims they make about dietary supplements. The FDA has the power to remove supplements from the market if it has sufficient scientific evidence to show the supplement is unsafe. Additionally, the FTC has the power to act against companies who make false and/or misleading marketing claims about a specific product. This includes acting against companies if the ingredients found in the supplement do not match label claims. While this does not ensure the safety of dietary supplements, it does provide a means for governmental oversight of the dietary supplement industry if adequate resources are provided to enforce DSHEA. Since inception of DSHEA, the FDA has required a number of supplement companies to submit evidence showing safety of their products and acted to remove a number of products sold as dietary supplements from sale in the U.S. due to safety concerns. Additionally, the FTC has acted against a number of supplement companies for misleading advertisements and/or structure and function claims.

As can be seen, although some argue that the dietary supplement industry is "unregulated" and/or may have suggestions for additional regulation, manufacturers of dietary supplements must adhere to a number of federal regulations before a product can go to market. Further, they must have evidence that the ingredients sold in their supplements are generally safe if requested to do so by the FDA. For this reason, over the last 10-15 years, most quality supplement companies have employed a team of researchers (many of whom are MS or PhD prepared exercise physiologists or sport nutrition specialists) who help educate the public about nutrition and exercise, provide input on product

development, conduct preliminary research on products, and/or assist in coordinating research trials conducted by independent research teams (e.g., university based researchers or clinical research sites). They also consult with marketing teams with the responsibility to ensure structure and function claims do not misrepresent results of research findings. This has increased job opportunities for sport nutrition specialists as well as enhanced opportunities for external funding for research groups interested in exercise nutrition research. While it is true that some companies use borrowed science, suppress negative findings, and/or exaggerate results from research studies, the trend in the nutrition industry is to develop scientifically sound supplements. This trend toward greater research support is the result of: 1.) attempts to honestly and accurately inform the public about results; 2.) efforts to have data to support safety and efficacy on products for FDA and the FTC; and/or, 3.) to provide scientific evidence to support advertising claims and increase sales. This trend is due in large part to greater scrutiny from the FDA and FTC as a result of increased consumer expectations and political pressure to ensure that companies sell quality products that have been shown to be safe and effective in clinical trials. In our experience, companies who adhere to these ethical standards prosper while those who do not struggle to adhere to FDA and FTC guidelines and lose consumer confidence. When this occurs, companies are often sued by consumers and/or are forced out of business because ultimately the consumer has the final word on whether a supplement or supplement company is credible or not.

HOW TO EVALUATE NUTRITIONAL ERGOGENIC AIDS

When you evaluate the ergogenic value of a nutritional supplement or training device/method, we recommend that you go through a process of evaluating the validity and scientific merit of claims made. This can be accomplished by evaluating the theoretical rationale behind the supplement/technique

and determining whether there is any well-controlled data showing the supplement/technique works. Training devices and supplements based on sound scientific rationale with supportive research showing effectiveness may be worth trying and/or recommending. However, those based on unsound scientific rationales and/or little to no data supporting the ergogenic value for people involved in intense training may not. The sport nutrition specialist should be a resource to help their clients interpret the scientific and medical research that may impact on their welfare and/or help them train more wisely. The following are the questions we recommend asking when evaluating the potential ergogenic value of a supplement.

Does The Theory Make Sense?

Most supplements that have been marketed to improve health and/or exercise performance are based on theoretical applications derived from basic and/or clinical research studies. Based on these preliminary studies, a training device or supplement is often marketed to people proclaiming the benefits observed in these basic research studies. Although the theory may sound good, critical analysis of the theory often reveals flaws in scientific logic and/or that the claims made don't quite match up with the literature cited. If you do your homework, you can discern whether a supplement has been based on sound scientific evidence or not. To do so, we suggest you read reviews about the training method, nutrient, and/or supplement from researchers who have been intimately involved in this line of research and/or consult reliable references about nutritional and herbal supplements⁴⁻⁸. We also suggest doing a search on the nutrient/supplement on the National Library of Medicine's Pub Med Online⁹. A quick look at these references will often help you know whether the theory is plausible or not. In our experience, proponents of ergogenic aids often overstate claims made about training devices and/or nutritional supplements while opponents of nutritional supplements and ergogenic aids

are either unaware and/or ignorant of research supporting their use. The sport nutrition specialist has the responsibility to know the literature and/or search available data bases to know whether there is merit or not to a proposed ergogenic aid.

Is There Any Scientific Evidence Supporting The Ergogenic Value?

The next question suggest asking is whether there is any well-controlled data showing the proposed ergogenic aid works as claimed in athletes or people involved in training. The first place we look is the list of references cited in marketing material supporting their claims. We look to see if the abstracts or articles cited are general references or specific studies that have evaluated the efficacy of the nutrient/supplement. We then critically evaluate the abstracts and articles by asking a series of questions.

- Are the studies simply basic research done in animals/clinical populations or have the studies been conducted on athletes? Studies reporting improved performance in rats may be insightful but research conducted on athletes is much more convincing.
- Were the studies well controlled? For ergogenic aid research, the study should be a placebo controlled, double blind, and randomized clinical trail if possible. This means that neither the researcher's nor the subject's were aware which group received the supplement or the placebo during the study and that the subjects were randomly assigned into the placebo or supplement group. At times, supplement claims have been based on poorly designed studies (i.e., small groups of subjects, no control group, use of unreliable tests, etc) and/or testimonials which may make interpretation much more difficult. Studies that are well controlled clinical trials provide stronger evidence as to the potential ergogenic value than those that are not well controlled.
- Do the studies report statistically significant results or are claims being made on non-significant means or trends reported? Appropriate statistical analysis of research results allows for an unbiased interpretation of data. Although studies reporting statistical trends may be of interest and lead researchers to conduct additional research, studies reporting statistically significant results are obviously more convincing. With this said, sport nutrition specialist must be careful not to commit type II statistical error (i.e., indicating that no differences were observed when a true effect was seen but not detected statistically). Since many studies on ergogenic aids (particularly in high level athletes) evaluate small numbers of subjects, results may not reach statistical significance even though large mean changes were observed. In these cases, additional research is warranted to further examine the potential ergogenic aid before conclusions can be made.
- Do the results of the studies cited match the claims made about the supplement? It is not unusual for marketing claims to greatly exaggerate the results found in the actual studies. Therefore, you should compare results observed in the studies to marketing claims. Reputable companies accurately report results of studies so that consumers can make informed decisions about whether to try a product or not.
- Were results of the study presented at a reputable scientific meeting and/or published in a peer-reviewed scientific journal? At times, claims are based on research that has either never been published or only published in an obscure journal. The best research is typically presented at respected scientific meetings and/or published in reputable peer-reviewed journals.
- Have the research findings been replicated at several different labs? The best way to know an ergogenic aid works is to see that results have been replicated in several studies preferably by a number of researchers. The most reliable ergogenic

aids are those in which a number of studies, conducted at different labs, have reported similar results.

Is The Supplement Legal And Safe?

The final question we ask is whether the supplement is legal and/or safe. Some athletic associations have banned the use of various nutritional supplements (e.g., prohormones, ephedra, etc). Obviously, if the supplement is banned, the sport nutrition specialist should discourage its use. In addition, many supplements have not been studied for long-term safety. People who consider taking nutritional supplements should be well aware of the potential side effects so that they can make an informed decision regarding whether to use a supplement or not. Additionally, they should consult with a knowledgeable physician to see if there are any underlying medical problems that may contraindicate use. When evaluating the safety of a supplement, we suggest looking to see if any side effects have been reported in the scientific or medical literature. In particular, we suggest determining how long a particular supplement has been studied, the dosages evaluated, and whether any side effects were observed. We also recommend consulting the PDR for nutritional supplements and herbal supplements to see if any side effects have been reported and/or there are any known drug interactions. If no side effects have been reported in the scientific/medical literature, we generally will view the supplement as safe for the length of time and dosages evaluated.

CLASSIFYING AND CATEGORIZING SUPPLEMENTS

Dietary supplements may contain carbohydrate, protein, fat, minerals, vitamins, herbs, and/or various plant/food extracts. Supplements can generally be classified as convenience supplements (e.g., energy bars, meal replacement powders, ready to drink supplements) designed to provide a convenient means of meeting caloric needs

and/or managing caloric intake, weight gain supplements, weight loss supplements, and performance enhancement supplements. Based on the above criteria, we generally categorize nutritional supplements into the following categories:

- I. **Apparently Effective.** Supplements that help people meet general caloric needs and/or the majority of research studies show is effective and safe.
- II. **Possibly Effective.** Supplements that initial studies support the theoretical rationale but that more research is needed to determine how the supplement may affect training and/or performance.
- III. **Too Early To Tell.** Supplements that the theory may make sense but there is insufficient research to support the use at this time.
- IV. **Apparently Ineffective.** Supplements that the theoretical rationale makes little scientific sense and/or research has clearly shown to be ineffective.

When a sport nutrition specialist counsels people who train, they should first evaluate their diet and training program. They should make sure that the athlete is eating an energy balanced, nutrient dense diet and that they are training intelligently. This is the foundation to build a good program. Following this, we recommend that they generally only recommend supplements in category I. If someone is interested in trying supplements in category II, they should make sure that they understand that these supplements are more experimental and that they may or may not see the type of results claimed. We recommend discouraging people from trying supplements in category III because there isn't enough data available on whether they work or not. However, if someone wants to try one of these supplements, they should understand that although there is some theoretical rationale, there is little evidence to support use at this time. Obviously, we do not support athletes taking supplements in categories IV. We believe that this approach is a more scientifically supportable and

balanced view than simply dismissing the use of all dietary supplements out of hand.

GENERAL DIETARY GUIDELINES FOR ACTIVE INDIVIDUALS

A well-designed diet that meets energy intake needs and incorporates proper timing of nutrients is the foundation upon which a good training program can be developed. Research has clearly shown that athletes that do not ingest enough calories and/or do not consume enough of the right type of macronutrients may impede training adaptations while athletes who consume a good diet can help the body adapt to training. Moreover, maintaining an energy deficient diet during training may lead to loss of muscle mass, increased susceptibility to illness, and increase prevalence of overreaching and/or overtraining. Incorporating good dietary practices as part of a training program is one way to help optimize training adaptations and prevent overtraining. The following overviews energy intake and major nutrient needs of active individuals.

Energy Intake

The first component to optimize training and performance through nutrition is to ensure the athlete is consuming enough calories to offset energy expenditure^{1, 10-12}. People who participate in a general fitness program (e.g., exercising 30 - 40 minutes per day, 3 times per week) can generally meet nutritional needs following a normal diet (e.g., 1,800 – 2,400 kcals/day or about 25 - 35 kcals/kg/day for a 50 – 80 kg individual) because their caloric demands from exercise are not too great (e.g., 200 – 400 kcals/session)¹. However, athletes involved in moderate levels of intense training (e.g., 2-3 hours per day of intense exercise performed 5-6 times per week) or high volume intense training (e.g., 3-6 hours per day of intense training in 1-2 workouts for 5-6 days per week) may expend 600 – 1,200 kcals or more per hour during exercise^{1, 13}. For this reason, their caloric

needs may approach 50 – 80 kcals/kg/day (2,500 – 8,000 kcals/day for a 50 – 100 kg athlete). For elite athletes, energy expenditure during heavy training or competition may be enormous. For example, energy expenditure for cyclists to compete in the Tour de France has been estimated as high as 12,000 kcals/day (150 - 200 kcals/kg/d for a 60 – 80 kg athlete)¹³⁻¹⁵. Additionally, caloric needs for large athletes (i.e., 100 – 150 kg) may range between 6,000 – 12,000 kcals/day depending on the volume and intensity of different training phases¹³.

Although some argue that athletes can meet caloric needs simply by consuming a well-balanced diet, it is often very difficult for larger athletes and/or athletes engaged in high volume/intense training to be able to eat enough food in order to meet caloric needs^{1, 11, 13-15}. Maintaining an energy deficient diet during training often leads to significant weight loss (including muscle mass), illness, onset of physical and psychological symptoms of overtraining, and reductions in performance¹². Nutritional analyses of athletes' diets have revealed that many are susceptible to maintaining negative energy intakes during training. Susceptible populations include runners, cyclists, swimmers, triathletes, gymnasts, skaters, dancers, wrestlers, boxers, and athletes attempting to lose weight too quickly¹¹. Additionally, female athletes have been reported to have a high incidence of eating disorders¹¹. Consequently, it is important for the sport nutrition specialist working with athletes to ensure that athletes are well-fed and consume enough calories to offset the increased energy demands of training and maintain body weight. Although this sounds relatively simple, intense training often suppresses appetite and/or alters hunger patterns so that many athletes do not feel like eating¹¹. Some athletes do not like to exercise within several hours after eating because of sensations of fullness and/or a predisposition to cause gastrointestinal distress. Further, travel and training schedules may limit food availability and/or

the types of food athletes are accustomed to eating. This means that care should be taken to plan meal times in concert with training as well as make sure athletes have sufficient availability of nutrient dense foods throughout the day for snacking between meals (e.g., drinks, fruit, carbohydrate/protein bars, etc)^{1, 10, 11}. For this reason, sport nutritionists' often recommend that athletes consume 4-6 meals per day and snack in between meals in order to meet energy needs. Use of nutrient dense energy bars and high calorie carbohydrate/protein supplements provides a convenient way for athletes to supplement their diet in order to maintain energy intake during training.

Carbohydrate

The second component to optimizing training and performance through nutrition is to ensure that athletes consume the proper amounts of carbohydrate, protein and fat in their diet. Individuals engaged in a general fitness program can typically meet macronutrient needs by consuming a normal diet (i.e., 45-55% carbohydrate [3-5 grams/kg/day], 10-15% protein [0.8 – 1.0 gram/kg/day], and 25-35% fat [0.5 – 1.5 grams/kg/day]). However, athletes involved in moderate and high volume training need greater amounts of carbohydrate and protein in their diet to meet macronutrient needs. For example, in terms of carbohydrate needs, athletes involved in moderate amounts of intense training (e.g., 2-3 hours per day of intense exercise performed 5-6 times per week) typically need to consume a diet consisting of 55-65% carbohydrate (i.e., 5-8 grams/kg/day or 250 – 1,200 grams/day for 50 – 150 kg athletes) in order to maintain liver and muscle glycogen stores^{1, 10}. Research has also shown that athletes involved in high volume intense training (e.g., 3-6 hours per day of intense training in 1-2 workouts for 5-6 days per week) may need to consume 8-10 grams/day of carbohydrate (i.e., 400 – 1,500 grams/day for 50 – 150 kg athletes) in order to maintain muscle glycogen levels^{1, 10}. This would be equivalent to

consuming 0.5 – 2.0 kg of spaghetti. Preferably, the majority of dietary carbohydrate should come from complex carbohydrates with a low to moderate glycemic index (e.g., grains, starches, fruit, maltodextrins, etc). However, since it is physically difficult to consume that much carbohydrate per day when an athlete is involved in intense training, many nutritionists and sport nutrition specialist recommend that athletes consume concentrated carbohydrate juices/drinks and/or consume high carbohydrate supplements to meet carbohydrate needs. While consuming this amount of carbohydrate is not necessary for the fitness minded individual who only trains 3-4 times per week for 30-60 minutes, it is essential for competitive athletes engaged in intense moderate to high volume training.

Protein

There has been considerable debate regarding protein needs of athletes¹⁶⁻²⁰. Initially, it was recommended that athletes do not need to ingest more than the RDA for protein (i.e., 0.8 to 1.0 g/kg/d for children, adolescents and adults). However, research over the last decade has indicated that athletes engaged in intense training need to ingest about 1.5 – 2 times the RDA of protein in their diet (1.5 to 2.0 g/kg/d) in order to maintain protein balance¹⁶⁻²⁰. If an insufficient amount of protein is obtained from the diet, an athlete will maintain a negative nitrogen balance which can increase protein catabolism and slow recovery. Over time, this may lead to lean muscle wasting and training intolerance^{1, 12}.

For people involved in a general fitness program, protein needs can generally be met by ingesting 0.8 – 1.0 grams/kg/day of protein. It is generally recommended that athletes involved in moderate amounts of intense training consume 1 – 1.5 grams/kg/day of protein (50 – 225 grams/day for a 50 – 150 kg athlete) while athletes involved in high volume intense

training consume 1.5 – 2.0 grams/kg/day of protein (75 – 300 grams/day for a 50 – 150 kg athlete)²¹. This protein need would be equivalent to ingesting 3 – 11 servings of chicken or fish per day for a 50 – 150 kg athlete²¹. Although smaller athletes typically can ingest this amount of protein in their normal diet, larger athletes often have difficulty consuming this much dietary protein. Additionally, a number of athletic populations have been reported to be susceptible to protein malnutrition (e.g., runners, cyclists, swimmers, triathletes, gymnasts, dancers, skaters, wrestlers, boxers, etc). Therefore, care should be taken to ensure that athletes consume a sufficient amount of quality protein in their diet in order to maintain nitrogen balance (e.g., 1.5 - 2 grams/kg/day).

However, it should be noted that not all protein is the same. Proteins differ based on the source that the protein was obtained, the amino acid profile of the protein, and the methods of processing or isolating the protein²². These differences influence availability of amino acids and peptides that have been reported to possess biological activity (e.g., α -lactalbumin, β -lactoglobulin, glycomacropptides, immunoglobulins, lactoperoxidases, lactoferrin, etc). Additionally, the rate and metabolic activity of the protein²². For example, different types of proteins (e.g., casein and whey) are digested at different rates which directly affect catabolism and anabolism²²⁻²⁵. Therefore, care should be taken not only to make sure the athlete consumes enough protein in their diet but also that the protein is high quality. The best dietary sources of low fat and high quality protein are light skinless chicken, fish, egg white and skim milk (casein and whey)²². The best sources of high quality protein found in nutritional supplements is whey, colostrum, casein, milk proteins and egg protein^{21,22}. Although some athletes may not need to supplement their diet with protein and some sport nutrition specialists may not think that protein supplements are

necessary, suggestions that it is unethical for a sport nutrition specialist to recommend that some athletes supplement their diet with protein in order to meet dietary protein needs and/or provide essential amino acids following exercise in order to optimize protein synthesis is clearly not supported by the literature.

Fat

The dietary recommendations of fat intake for athletes are similar to or slightly greater than those recommended for non-athletes in order to promote health. Maintenance of energy balance, replenishment of intramuscular triacylglycerol stores and adequate consumption of essential fatty acids are of greater importance among athletes and allow for somewhat increased intake²⁶. This depends on the athlete's training state and goals. For example, higher-fat diets appear to maintain circulating testosterone concentrations better than low-fat diets²⁷⁻²⁹. This has relevance to the documented testosterone suppression which can occur during volume-type overtraining³⁰. Generally, it is recommended that athletes consume a moderate amount of fat (approximately 30% of their daily caloric intake), while increases up to 50% of kcal can be safely ingested by athletes during regular high-volume training²⁶. For athletes attempting to decrease body fat, however, it has been recommended that they consume 0.5 to 1 g/kg/d of fat¹. The reason for this is that some weight loss studies indicate that people who are most successful in losing weight and maintaining the weight loss are those who ingest less than 40 g/d of fat in their diet^{31,32} although this is not always the case³³. Certainly, the type of dietary fat (e.g. n-6 versus n-3; saturation state) is a factor in such research and could play an important role in any discrepancies^{34,35}. Strategies to help athletes manage dietary fat intake include teaching them which foods contain various types of fat so that they can make better

food choices and how to count fat grams^{1, 11}.

Strategic Eating and Refueling

In addition to the general nutritional guidelines described above, research has also demonstrated that timing and composition of meals consumed may play a role in optimizing performance, training adaptations, and preventing overtraining^{1, 10, 36, 37}. In this regard, it takes about 4 hours for carbohydrate to be digested and begin to be stored as muscle and liver glycogen. Consequently, pre-exercise meals should be consumed about 4 to 6 h before exercise¹⁰. This means that if an athlete trains in the afternoon, breakfast is the most important meal to top off muscle and liver glycogen levels. Research has also indicated that ingesting a light carbohydrate and protein snack 30 to 60 min prior to exercise (e.g., 50 g of carbohydrate and 5 to 10 g of protein) serves to increase carbohydrate availability toward the end of an intense exercise bout^{38, 39}. This also serves to increase availability of amino acids and decrease exercise-induced catabolism of protein^{36, 38, 39}.

When exercise lasts more than one hour, athletes should ingest glucose/electrolyte solution (GES) drinks in order to maintain blood glucose levels, help prevent dehydration, and reduce the immunosuppressive effects of intense exercise^{10, 40-45}. Following intense exercise, athletes should consume carbohydrate and protein (e.g., 1 g/kg of carbohydrate and 0.5 g/kg of protein) within 30 min after exercise as well as consume a high carbohydrate meal within two hours following exercise^{1, 36, 37}. This nutritional strategy has been found to accelerate glycogen resynthesis as well as promote a more anabolic hormonal profile that may hasten recovery⁴⁶⁻⁴⁸. Finally, for 2 to 3 days prior to competition, athletes should taper training by 30 to 50% and consume 200 to 300 g/d of extra carbohydrate in their diet. This carbohydrate loading technique

has been shown to supersaturate carbohydrate stores prior to competition and improve endurance exercise capacity^{1, 10, 37}. Thus, the type of meal and timing of eating are important factors in maintaining carbohydrate availability during training and potentially decreasing the incidence of overtraining.

Vitamins

Vitamins are essential organic compounds which serve to regulate metabolic processes, energy synthesis, neurological processes, and prevent destruction of cells. There are two primary classifications of vitamins: fat and water soluble. The fat soluble vitamins include vitamins A, D, E, & K. The body stores fat soluble vitamins and therefore excessive intake may result in toxicity. Water soluble vitamins are B vitamins and vitamin C. Since these vitamins are water soluble, excessive intake of these vitamins are eliminated in urine. Table 1 describes RDA, proposed ergogenic benefit, and summary of research findings for fat and water soluble vitamins. Although research has demonstrated that specific vitamins may possess some health benefit (e.g., vitamin E, niacin, folic acid, vitamin C, etc), few have been reported to directly provide ergogenic value for athletes. However, some vitamins may help athletes tolerate training to a better degree by reducing oxidative damage (vitamin E, C) and/or help to maintain a healthy immune system during heavy training (vitamin C). Theoretically, this may help athletes tolerate heavy training leading to improved performance. The remaining vitamins reviewed appear to have little ergogenic value for athletes who consume a normal, nutrient dense diet. Since dietary analyses of athletes have found deficiencies in caloric and vitamin intake, many sport nutritionists' recommend that athletes consume a low-dose one a day multivitamin and/or a vitamin enriched post-workout carbohydrate/protein supplement during periods of heavy training. The American Medical Association also recently evaluated

the available medical literature and recommended that Americans consume a one-a-day low-dose multivitamin in order to promote general health. Suggestions that there is no benefit of vitamin supplementation for athletes and/or it is unethical for a sport nutrition specialist to recommend that their clients take a one-a-day multi-vitamin and/or suggest taking other vitamins that may reduce cholesterol levels (niacin), serve as antioxidants (Vitamin E), decrease risk to heart disease (niacin, Vitamin E), or may help maintain a health immune system (Vitamin C) is not consistent with current available literature.

Minerals

Minerals are essential inorganic elements necessary for a host of metabolic processes. Minerals serve as structure for tissue, important components of enzymes and hormones, and regulators of metabolic and neural control. Some minerals have been found to be deficient in athletes or become deficient in response to training and/or prolonged exercise. When mineral status is inadequate, exercise capacity may be reduced. Dietary supplementation of minerals in deficient athletes has generally been found to improve exercise capacity. Additionally, supplementation of specific minerals in non-deficient athletes has also been reported to affect exercise capacity. Table 2 describes minerals that have been purported to affect exercise capacity in athletes. Of the minerals reviewed, several appear to possess health and/or ergogenic value for athletes under certain conditions. For example, calcium supplementation in athletes susceptible to premature osteoporosis may help maintain bone mass. There is also recent evidence that dietary calcium may help manage body composition. Iron supplementation in athletes prone to iron deficiencies and/or anemia has been reported to improve exercise capacity. Sodium phosphate loading has been reported to increase maximal oxygen uptake, anaerobic threshold, and improve endurance exercise capacity by 8 to 10%. Increasing dietary

availability of salt (sodium chloride) during the initial days of exercise training in the heat has been reported to help maintain fluid balance and prevent dehydration. Finally, zinc supplementation during training has been reported to decrease exercise-induced changes in immune function. Consequently, somewhat in contrast to vitamins, there appear to be several minerals that may enhance exercise capacity and/or training adaptations for athletes under certain conditions. However, although ergogenic value has been purported for remaining minerals, there is little evidence that boron, chromium, magnesium, or vanadium affect exercise capacity or training adaptations in healthy individuals eating a normal diet. Suggestions that there is no benefit of mineral supplementation for athletes and/or it is unethical for a sport nutrition specialist to recommend that their clients take minerals that research has shown may affect health and/or performance is not consistent with current available literature.

Water

The most important nutritional ergogenic aid for athletes is water. Exercise performance can be significantly impaired when 2% or more of body weight is lost through sweat. For example, when a 70-kg athlete loses more than 1.4 kg of body weight during exercise (2%), performance capacity is often significantly decreased. Further, weight loss of more than 4% of body weight during exercise may lead to heat illness, heat exhaustion, heat stroke, and possibly death⁴⁵. For this reason, it is critical that athletes consume a sufficient amount of water and/or GES sports drinks during exercise in order to maintain hydration status. The normal sweat rate of athletes ranges from 0.5 to 2.0 L/h depending on temperature, humidity, exercise intensity, and their sweat response to exercise⁴⁵. This means that in order to maintain fluid balance and prevent dehydration, athletes need to ingest 0.5 to 2 L/h of fluid in order to offset weight loss. This requires frequent ingestion of 6-8 oz of

cold water or a GES sports drink every 5 to 15-min during exercise^{45, 49-52}. Athletes and should not depend on thirst to prompt them to drink because people do not typically get thirsty until they have lost a significant amount of fluid through sweat. Additionally, athletes should weigh themselves prior to and following exercise training to ensure that they maintain proper hydration^{45, 49-52}. The athlete should consume 3 cups of water for every pound lost during exercise in order adequately rehydrate themselves⁴⁵. Athletes should train themselves to tolerate drinking greater amounts of water during training and make sure that they consume more fluid in hotter/humid environments. Preventing dehydration during exercise is one of the most effective ways to maintain exercise capacity. Finally, inappropriate and excessive weight loss techniques (e.g., cutting weight in saunas, wearing rubber suits, severe dieting, vomiting, using diuretics, etc) are extremely dangerous and should be prohibited. Sport nutrition specialists can play an important role in educating athletes and coaches about proper hydration methods and supervising fluid intake during training and competition.

DIETARY SUPPLEMENTS AND ATHLETES

Most of the work we do with athletes regarding sport nutrition is to teach them and their coaches how to structure their diet and time food intake to optimize performance and recovery. Dietary supplements can play a meaningful role in helping athletes consume the proper amount of calories, carbohydrate, and protein in their diet. However, they should be viewed as supplements to the diet, not replacements for a good diet. While it is true that most dietary supplements available for athletes have little scientific data supporting their potential role to enhance training and/or performance, it is also true that a number of nutrients and/or dietary supplements have been shown to help improve performance and/or recovery. This can help augment the normal diet to help optimize performance. Sport nutrition

specialists must be aware of the current data regarding nutrition, exercise, and performance and be honest about educating their clients about results of various studies (whether pro or con). With the proliferation of information available about nutritional supplements to the consumer, the sport nutrition specialist, nutritionist, and nutrition industry lose credibility when they do not accurately describe results of various studies to the public. The following overviews several classifications of nutritional supplements that are often taken by athletes and categorizes them into apparently effective, possibly effective, too early to tell, and apparently ineffective supplements based on my interpretation of the literature. It should be noted that this analysis will primarily focus on whether the proposed nutrient has been found to affect exercise and/or training adaptations based on the current available literature. Additional research may reveal it may or may not possess ergogenic value which may then change its classification. It should be also noted that although there may be little ergogenic value to some nutrients, there may be some potential health benefits that may be helpful for some populations. Therefore, just because a nutrient does not appear to affect performance and/or training adaptations, that does not mean it may not have possible health benefits.

Convenience Supplements

Convenience supplements are meal replacement powders (MRP's), ready to drink supplements (RTD's), energy bars, and energy gels. They currently represent the largest segment of nutrition industry representing 50 – 75% of most company's sales. They are typically fortified with 33 – 50% of the RDA for vitamins and minerals and typically differ on the amount of carbohydrate, protein, and fat they contain. They may also differ based whether they are fortified with various nutrients purported to promote weight gain, enhance weight loss, and/or improve performance. Most people view these supplements as a high quality

snacks and/or use them to help control caloric intake when trying to gain and/or lose weight. In our view, MRP's, RTD's, and energy bars/gels can provide a convenient way for people to meet specific dietary needs and/or serve as good alternatives to fast food. Use of these types of products can be particularly helpful in providing carbohydrate, protein, and other nutrients prior to and/or following exercise in an attempt to optimize nutrient intake when an athlete doesn't have time to sit down for a good meal. However, they should be used to improve dietary availability of macronutrients – not as a replacement for a good diet. Care should also be taken to make sure they do not contain any banned or prohibited nutrients.

Muscle Building Supplements

The following provides an analysis of the literature regarding purported weight gain supplements and our general interpretation of how they should be categorized based on this information. Table 3 summarizes how we currently classify the ergogenic value of a number of purported performance-enhancing, muscle building, and fat loss supplements based on an analysis of the available scientific evidence.

Apparently Effective

Weight Gain Powders. One of the most common means athletes have employed to increase muscle mass is to add extra calories to the diet. Most athletes “bulk up” in this manner by consuming extra food and/or weight gain powders. Studies have consistently shown that simply adding an extra 500 – 1,000 calories per day to your diet will promote weight gain^{20,36}. However, only about 30 – 50% of the weight gained on high calorie diets is muscle while the remaining amount of weight gained is fat. Consequently, increasing muscle mass by ingesting a high calorie can help you build muscle but the accompanying increase in body fat may not be desirable for everyone.

Therefore, we typically do not recommend this type of weight gain approach.

Creatine. In our view, the most effective nutritional supplement available to athletes to increase high intensity exercise capacity and muscle mass during training is creatine. Numerous studies have indicated that creatine supplementation increases body mass and/or muscle mass during training⁵³. Gains are typically 2 – 5 pounds greater than controls during 4 – 12 weeks of training⁵⁴. The gains in muscle mass appear to be a result of an improved ability to perform high intensity exercise enabling an athlete to train harder and thereby promote greater training adaptations and muscle hypertrophy⁵⁵⁻⁵⁷. The only clinically significant side effect reported from creatine supplementation has been weight gain^{36, 53, 54, 58}. Although concerns have been raised about the safety and possible side effects of creatine supplementation^{59, 60}, recent long-term safety studies have reported no apparent side effects^{58, 61, 62} and/or that creatine may lessen the incidence of injury during training⁶³⁻⁶⁵. Consequently, supplementing the diet with creatine and/or creatine containing formulations seems to be a safe and effective method to increase muscle mass.

β-hydroxy β-methylbutyrate (HMB). HMB is a metabolite of the amino acid leucine. Leucine and metabolites of leucine have been reported to inhibit protein degradation⁶⁶. Supplementing the diet with 1.5 to 3 g/d of calcium HMB has been typically reported to increase muscle mass and strength particularly among untrained subjects initiating training⁶⁷⁻⁷² and the elderly⁷³. Gains in muscle mass are typically 0.5 to 1 kg greater than controls during 3 – 6 weeks of training. There is also recent evidence that HMB may lessen the catabolic effects of prolonged exercise⁷⁴ and that there may be additive effects of co-ingesting HMB with creatine^{75, 76}. However, the effects of HMB supplementation in athletes are less clear. Most studies conducted on trained subjects have reported non-significant gains in muscle

mass possibly due to a greater variability in response of HMB supplementation among athletes⁷⁷⁻⁷⁹. Consequently, there is fairly good evidence showing that HMB may enhance training adaptations in individuals initiating training. However, additional research is necessary to determine whether HMB may enhance training adaptations in athletes.

Possibly Effective

Branched Chain Amino Acids (BCAA).

BCAA supplementation has been reported to decrease exercise-induced protein degradation and/or muscle enzyme release (an indicator of muscle damage) possibly by promoting an anti-catabolic hormonal profile^{36, 38, 80}.

Theoretically, BCAA supplementation during intense training may help minimize protein degradation and thereby lead to greater gains in fat-free mass. There is some evidence to support this hypothesis. For example, Schena and colleagues⁸¹ reported that BCAA supplementation (~10 g/d) during 21-days of trekking at altitude increased fat free mass (1.5%) while subjects ingesting a placebo had no change in muscle mass. Bigard and associates⁸² reported that BCAA supplementation appeared to minimize loss of muscle mass in subjects training at altitude for 6-weeks. Finally, Candeloro and coworkers⁸³ reported that 30 days of BCAA supplementation (14 grams/day) promoted a significant increase in muscle mass (1.3%) and grip strength (+8.1%) in untrained subjects. Although more research is necessary, these findings suggest that BCAA supplementation may have some impact on body composition.

Essential Amino Acids (EAA). Recent studies have indicated that ingesting 3 to 6 g of EAA prior to^{84, 85} and or following exercise stimulates protein synthesis⁸⁵⁻⁹². Theoretically, this may enhance gains in muscle mass during training. To support this theory, a recent study by Esmarck and colleagues⁹³ found that ingesting EAA with carbohydrate immediately following

resistance exercise promoted significantly greater training adaptations as compared to waiting until 2-hours after exercise to consume the supplement. Although more data is needed, there appears to be strong theoretical rationale and some supportive evidence that EAA supplementation may enhance protein synthesis and training adaptations.

Glutamine. Glutamine is the most plentiful non-essential amino acid in the body and plays a number of important physiological roles³⁶. Glutamine has been reported to increase cell volume and stimulate protein⁹⁴⁻⁹⁶ and glycogen synthesis⁹⁷. Theoretically, glutamine supplementation prior to and/or following exercise (e.g., 6-10 g) may help to optimize cell hydration and protein synthesis during training leading to greater gains in muscle mass and strength^{36, 98}. In support of this hypothesis, a recent study by Colker and associates⁹⁹ found that subjects who supplemented their diet with glutamine (5 grams) and BCAA (3 grams) enriched whey protein during training promoted about a 2 pound greater gain in muscle mass and greater gains in strength than ingesting whey protein alone. Although more data is needed, there appears to be a strong scientific rationale and some preliminary evidence to indicate that glutamine may help build muscle.

Protein. As previously described, research has indicated that people undergoing intense training may need additional protein in their diet to meet protein needs (i.e., 1.5 – 2.0 grams/day). People who do not ingest enough protein in their diet may slow recovery and training adaptations³⁶. Protein supplements offer a convenient way to ensure that athletes consume quality protein in the diet and meet their protein needs. However, ingesting additional protein beyond that necessary to meet protein needs does not appear to promote additional gains in strength and muscle mass. The research focus over recent years has been to determine whether different types of protein (e.g., whey, casein, soy, milk proteins, colostrum, etc) and/or various

biologically active protein subtypes and peptides (e.g., α -lactalbumin, β -lactoglobulin, glycomacropeptides, immunoglobulins, lactoperoxidases, lactoferrin, etc) have varying effects on the physiological, hormonal, and/or immunological responses to training. In addition, whether timing of protein intake may play a role in protein synthesis and training adaptations⁸⁵⁻⁹². Although more research is necessary in this area, research clearly indicates that protein needs of individuals engaged in intense training are elevated, that different types of protein have varying effects on anabolism and catabolism, that different types of protein subtypes and peptides have unique physiological effects, and that timing of protein intake may play an important role in optimizing protein synthesis following exercise. Therefore, it is simplistic and misleading to suggest that there is no data supporting contentions that athletes need more protein in their diet and/or there is no potential ergogenic value of incorporating different types of protein into the diet.

Too Early to Tell

α -ketoglutarate (α -KG). α -KG is an intermediate in the Krebs cycle that is involved in aerobic energy metabolism. There is some clinical evidence that α -KG may serve as an anticatabolic nutrient after surgery^{100, 101}. However, it is unclear whether α -KG supplementation during training may affect training adaptations.

α -Ketoisocaproate (KIC). KIC is a branched-chain keto acid that is a metabolite of leucine metabolism. In a similar manner as HMB, leucine and metabolites of leucine are believed to possess anticatabolic properties¹⁰². There is some clinical evidence that KIC may spare protein degradation in clinical populations^{103, 104}. Theoretically, KIC may help minimize protein degradation during training possibly leading to greater training adaptations. However, we are not aware of any studies that have evaluated the effects of KIC supplementation during training on body

composition.

Ecdysterones. Ecdysterones (also known as ectysterone, 20 Beta-Hydroxyecdysterone, turkesterone, ponasterone, ecdysone, or ecdystene) are naturally derived phytoecdysteroids (i.e., insect hormones). They are typically extracted from the herbs *Leuzea rhaponticum* sp., *Rhaponticum carthamoides*, or *Cyanotis vaga*. They can also be found in high concentrations in the herb *Suma* (also known as Brazilian Ginseng or *Pfaffia*). Research from Russia and Czechoslovakia conducted over the last 30 years indicates that ecdysterones may possess some potentially beneficial physiological effects in insects and animals¹⁰⁵⁻¹⁰⁹. However, since most of the data on ecdysterones have been published in obscure journals, results are difficult to interpret. While future studies may find some ergogenic value of ecdysterones, it is our view that it is too early to tell whether phytoecdysteroids serve as a safe and effective nutritional supplement for athletes.

Growth Hormone Releasing Peptides (GHRP) and Secretagogues. Research has indicated that growth hormone releasing peptides (GHRP) and other non-peptide compounds (secretagogues) appear to help regulate growth hormone (GH) release^{110, 111}. These observations have served as the basis for development of nutritionally-based GH stimulators (e.g., amino acids, pituitary peptides, "pituitary substances", macuna pruriens, broad bean, alpha GPC, etc). Although there is clinical evidence that pharmaceutical grade GHRP's and some non-peptide secretagogues can increase GH and IGF-1 levels at rest and in response to exercise, it is currently unknown whether any of these nutritional alternatives would increase GH and/or affect training adaptations.

Isoflavones. Isoflavones are naturally occurring non-steroidal phytoestrogens that have a similar chemical structure as the ipriflavone (a synthetic flavonoid drug used in

the treatment of osteoporosis)¹¹²⁻¹¹⁴. For this reason, soy protein (which is an excellent source of isoflavones) and isoflavone extracts have been investigated in the possible treatment of osteoporosis. Results of these studies have shown promise in preventing declines in bone mass in post-menopausal women as well as reducing risks to side effects associated with estrogen replacement therapy. More recently, the isoflavone extracts 7-isopropoxyisoflavone (ipriflavone) and 5-methyl-7-methoxy-isoflavone (methoxyisoflavone) have been marketed as “powerful anabolic” substances. These claims have been based on research described in patents filed in Hungary in the early 1970s^{115, 116}. Although the data presented in the patents are interesting, there is currently no peer-reviewed data indicating that isoflavone supplementation affects exercise, body composition, or training adaptations.

Ornithine- α -ketoglutarate (OKG). OKG is another nutrient believed to possess anabolic/catabolic effect. Animal and clinical studies have suggested that patients administered OKG experienced improved protein balance^{115, 116}. Theoretically, OKG may provide some value for athletes engaged in intense training. A recent study by Chetlin and colleagues¹¹⁷ reported that OKG supplementation (10 grams/day) during 6-weeks of resistance training promoted greater gains in bench press. However, no significant differences were observed in squat strength, training volume, gains in muscle mass, or fasting insulin and growth hormone. Therefore, additional research is needed before conclusions can be drawn.

Sulfo-Polysaccharides (Myostatin Inhibitors). Myostatin or growth differentiation factor 8 (GDF-8) is a transforming growth factor that has been shown to serve as a genetic determinant of the upper limit of muscle size and growth¹¹⁸. Recent research has indicated that eliminating and/or inhibiting myostatin gene expression in mice¹¹⁹ and cattle¹²⁰⁻¹²² promotes marked increases in muscle mass during early growth

and development. The result is that these animals experience what has been termed as a “double-muscle” phenomenon apparently by allowing muscle to grow beyond its normal genetic limit. In agriculture research, eliminating and/or inhibiting myostatin may serve as an effective way to optimize animal growth leading to larger, leaner, and a more profitable livestock yield. In humans, inhibiting myostatin gene expression has been theorized as a way to prevent or slow down muscle wasting in various diseases, speed up recovery of injured muscles, and/or promote increases in muscle mass and strength in athletes¹²³. While these theoretical possibilities may have great promise, research on the role of myostatin inhibition on muscle growth and repair is in the very early stages – particularly in humans. There is some evidence that myostatin levels are higher in the blood of HIV positive patients who have experience muscle wasting and that myostatin levels negatively correlate with muscle mass¹¹⁸. There is also evidence that myostatin gene expression may be fiber specific and that myostatin levels may be influenced by immobilization in animals¹²⁴. Additionally, a recent study by Ivey and colleagues¹²³ reported that female athletes with a less common myostatin allele (a genetic subtype that may be more resistant to myostatin) experienced greater gains in muscle mass during training and less loss of muscle mass during detraining. No such pattern was observed in men with varying amounts of training histories and muscle mass. These early studies suggest that myostatin may play a role in regulating muscle growth to some degree. Recently, some nutrition supplement companies have marketed sulfo-polysaccharides (derived from a sea algae called *Cytoseira canariensis*) as a way to partially bind the myostatin protein in serum. Although this theory is interesting and studies examining this hypothesis are underway, there is currently no published data supporting the use of sulfo-polysaccharides as a muscle building supplement.

Smilax Officinalis (SO). SO is a compound which contains plant sterols purported to enhance immunity as well as provide an androgenic effect on muscle growth¹. Some data supports the potential immune enhancing effects of SO. However, we are not aware of any data that show that SO supplementation increases muscle mass during training.

Zinc/Magnesium Aspartate (ZMA). ZMA formulations have recently become a popular supplement purported to promote anabolism at night. The theory is based on studies suggesting that zinc and magnesium deficiency may reduce the production of testosterone and insulin like growth factor (IGF-1). ZMA supplementation has been theorized to increase testosterone and IGF-1 leading to greater recovery, anabolism, and strength during training. In support of this theory, Brilla and Conte¹²⁵ reported that a zinc-magnesium formulation increased testosterone and IGF-1 (two anabolic hormones) leading to greater gains in strength in football players participating in spring training. While these data are interesting, more research is needed to further evaluate the role of ZMA on body composition and strength during training before conclusions can be drawn.

Apparently Ineffective

Boron. Boron is a trace mineral proposed to increase testosterone levels and promote anabolism. Several studies have evaluated the effects of boron supplementation during training on strength and body composition alterations. These studies indicate that boron supplement (2.5 mg/d) appears to have no impact on muscle mass or strength^{126, 127}.

Chromium. Chromium is a trace mineral that is involved in carbohydrate and fat metabolism. Clinical studies have suggested that chromium may enhance the effects of insulin particularly in diabetic populations. Since insulin is an anti-catabolic hormone and has been reported to affect protein synthesis, chromium supplementation has been

theorized to serve as an anabolic nutrient. Theoretically, this may increase anabolic responses to exercise. Although some initial studies reported that chromium supplementation increased gains in muscle mass and strength during training particularly in women¹²⁸⁻¹³⁰, most well-controlled that have been conducted since then have reported no benefit in healthy individuals taking chromium (200-800 mcg/d) for 4 to 16-weeks during training¹³¹⁻¹³⁷. Consequently, it appears that although chromium supplementation may have some therapeutic benefits for diabetics, chromium does not appear to be a muscle-building nutrient for athletes.

Conjugated Linoleic Acids (CLA). Animal studies indicate that adding CLA to dietary feed decreases body fat, increases muscle and bone mass, has anti-cancer properties, enhances immunity, and inhibits progression of heart disease¹³⁸⁻¹⁴⁰. Consequently, CLA supplementation in humans has been suggested to help manage body composition, delay loss of bone, and provide health benefit. Although animal studies are impressive¹⁴¹⁻¹⁴³ and a some studies suggests benefit at some but not all dosages^{144, 145}, most studies conducted on humans show little to no effect on body composition or muscle growth.^{146, 147}

Gamma Oryzanol (Ferulic Acid). Gamma oryzanol is a plant sterol theorized to increase anabolic hormonal responses during training¹⁴⁸. Although data are limited, one study reported no effect of 0.5 g/d of gamma oryzanol supplementation on strength, muscle mass, or anabolic hormonal profiles during 9-weeks of training¹⁴⁹.

Anabolic Steroids & Prohormones

Testosterone and growth hormone are two primary hormones in the body that serve to promote gains in muscle mass (i.e., anabolism) and strength while decreasing muscle breakdown (catabolism) and fat mass¹⁵⁰⁻¹⁵⁴. Testosterone also promotes male sex characteristics (e.g., hair, deep voice, etc)¹⁵⁴. Low level anabolic steroids are often

prescribed by physicians to prevent loss of muscle mass for people with various diseases and illnesses¹⁵⁵⁻¹⁶⁶. It is well known that athletes have experimented with large doses of anabolic steroids in an attempt to enhance training adaptations, increase muscle mass, and/or promote recovery during intense training¹⁵⁰⁻¹⁵⁴. Research has generally shown that use of anabolic steroids and growth hormone during training can promote gains in strength and muscle mass^{150, 160, 167-174}. However, a number of potentially life threatening adverse effects of steroid abuse have been reported including liver and hormonal dysfunction, hyperlipidemia (high cholesterol), increased risk to cardiovascular disease, and behavioral changes (i.e., steroid rage)^{169, 175-179}. Some of the adverse effects associated with the use of these agents are irreversible, particularly in women¹⁷⁶. For this reason, anabolic steroids have been banned by most sport organizations and should be avoided unless prescribed by a physician to treat an illness.

Prohormones (androstenedione, 4-androstenediol, 19-nor-4-androstenedione, 19-nor-4-androstenediol, 7-keto DHEA, and DHEA, etc) are naturally derived precursors to testosterone or other anabolic steroids. Prohormones have become popular among body builders because they believe they are natural boosters of anabolic hormones. Consequently, a number of over-the-counter supplements contain prohormones. While there is a strong theoretical rationale that prohormones may increase testosterone levels, there is virtually no evidence that these compounds affect training adaptations in younger men with normal hormone levels. In fact, most studies indicate that they do not affect testosterone and that some may actually increase estrogen levels and reduce HDL-cholesterol^{169, 180-186}. Consequently, although there may be some potential applications for older individuals to replace diminishing androgen levels, it appears that prohormones have no training value. Since prohormones are "steroid-like compounds", most athletic organizations have banned their use. Use of

nutritional supplements containing prohormones will result in a positive drug test for anabolic steroids. Use of supplements knowingly or unknowingly containing prohormones have been believed to have contributed to a number of recent positive drug tests among athletes. Consequently, care should be taken to make sure that any supplement an athlete considers taking does not contain prohormone precursors particularly if their sport bans and tests for use of such compounds.

Tribulus Terrestris. Tribulus terrestris (also known as puncture weed/vine or caltrops) is a plant extract that has been suggested to stimulate leutinizing hormone (LH) which stimulates the natural production of testosterone¹⁰². Consequently, Tribulus has been marketed as a supplement that can increase testosterone and promote greater gains in strength and muscle mass during training. Several recent studies have indicated that Tribulus supplementation appears to have no effects on body composition or strength during training^{187, 188}.

Vanadyl Sulfate (Vanadium). In a similar manner as chromium, vanadyl sulfate is a trace mineral that has been found to affect insulin-sensitivity and may affect protein and glucose metabolism¹⁰². For this reason, vanadyl sulfate has been purported to increase muscle mass and strength during training. Although there may be some clinical benefits for diabetics, vanadyl sulfate supplementation does not appear to have any effect on strength or muscle mass during training in non-diabetic individuals^{189, 190}.

Weight Loss Supplements

Although exercise and proper diet remain the best way to promote weight loss and/or manage body composition, a number of nutritional approaches have been investigated as possible weight loss methods (with or without exercise). The following overviews the major types of weight loss products available and discusses whether any available

research supports their use. See Table 3 for a summary.

Apparently Effective

Low Calorie Diet Foods & Supplements.

Most of the products in this category represent low fat/carbohydrate, high protein food alternatives¹⁹¹. They typically consist of pre-packaged food, bars, MRP, or RTD supplements. They are designed to provide convenient foods/snacks to help people follow a particular low calorie diet plan. In the scientific literature, diets that provide less than 1000 calories per day are known as very low calorie diets (VLCD's). Pre-packaged food, MRP's, and/or RTD's are often provided in VLCD plans to help people cut calories. In most cases, VLCD plans recommend behavioral modification and that people start a general exercise program.

Research on the safety and efficacy of people maintaining VLCD's generally indicate that they can promote weight loss. For example, Hoie et al¹⁹² reported that maintaining a VLCD for 8-weeks promoted a 27 lbs (12.6%) loss in total body mass, a 21 lbs loss in body fat (23.8%), and a 7 lbs (5.2%) loss in lean body mass in 127 overweight volunteers. Bryner and colleagues¹⁹³ reported that addition of a resistance training program while maintaining a VLCD (800 kcal/d for 12-weeks) resulted in a better preservation of lean body mass and resting metabolic rate compared to subjects maintaining a VLCD while engaged in an endurance training program. Kern and coworkers¹⁹⁴ reported that a medically supervised weight loss program involving behavioral modification and VLCD promoted a 51 lbs weight loss and that 61% of subjects maintained at least 50% of the weight loss at 12 and 18 months follow-up. Recent studies indicate that high protein/low fat VLCD's may be better than high carbohydrate/low fat diets in promoting weight loss^{32, 195-198}. The reason for this is that typically when people lose weight about 40-50% of the weight loss is muscle which decreases resting energy expenditure.

Increasing protein intake during weight loss helps preserve muscle mass and resting energy expenditure to a better degree than high carbohydrate diets¹⁹⁹. These findings and others indicate that VLCD's (typically using MRP's and/or RTD's as a means to control caloric intake) can be effective particularly as part of an exercise and behavioral modification program. Most people appear to maintain at least half of the initial weight lost for 1-2 years but tend to regain most of the weight back within 2-5 years. Therefore, although these diets may help people lose weight on the short-term, it is essential people who use them follow good diet and exercise practices in order to maintain the weight loss.

Ephedra, Caffeine, and Silicin.

Thermogenics are supplements designed to stimulate metabolism thereby increasing energy expenditure and promote weight loss. They typically contain the "ECA" stack of ephedra alkaloids (e.g., Ma Huang, 1R,2S Nor-ephedrine HCl, Sida Cordifolia), caffeine (e.g., Gaurana, Bissey Nut, Kola) and aspirin/salicin (e.g., Willow Bark Extract). More recently, other potentially thermogenic nutrients have been added to various thermogenic formulations. For example, thermogenic supplements may also contain synephrine (e.g., Citrus Aurantium, Bitter Orange), calcium & sodium phosphate, thyroid stimulators (e.g., guggulsterones, L-tyrosine, iodine), cayenne & black pepper, and ginger root.

A significant amount of research has evaluated the safety and efficacy of EC and ECA type supplements. Studies show that use of synthetic or herbal sources of ephedrine and caffeine (EC) promote about 2 lbs of extra weight loss per month while dieting (with or without exercise) and that EC supplementation is generally well tolerated in healthy individuals²⁰⁰⁻²⁰⁹. For example, Boozer et al²⁰¹ reported that 8-weeks of ephedrine (72 mg/d) and caffeine (240 mg/d) supplementation promoted a 9 lbs loss in body mass and a 2.1 % loss in body fat with

minor side effects. Molnar and associates²⁰⁰ reported that overweight children treated for 20 weeks with ephedrine and caffeine observed a 14.4% loss in body mass and a 6.6% decrease in body fat with no differences in side effects. Interestingly, Greenway and colleagues²⁰⁸ reported that EC supplementation was a more cost-effective treatment for reducing weight, cardiac risk, and LDL cholesterol than several weight loss drugs (fenfluramine with mazindol or phentermine). Finally, Boozer and associates²⁰⁰ reported that 6-months of herbal EC supplementation promoted weight loss with no clinically significant adverse effects in healthy overweight adults. Less is known about the safety and efficacy of synephrine, thyroid stimulators, cayenne/black pepper and ginger root.

Despite these findings, the Food and Drug Administration (FDA) has recently banned the sale of ephedra containing supplements. The rationale has been based on reports to adverse event monitoring systems and in the media suggesting a link between intake of ephedra and a number of severe medical complications (e.g., high blood pressure, elevated heart rate, arrhythmias, sudden death, heat stroke, etc)^{210, 211}. Although results of available clinical studies do not show these types of adverse events and the ban is in the process of being contested legally, ephedra is no longer available as an ingredient in dietary supplements. Consequently, thermogenic supplements now contain other nutrients believed to increase energy expenditure (e.g., synephrine, green tea, etc). Anyone contemplating taking thermogenic supplements should carefully consider the potential side effects, discuss possible use with a knowledgeable physician, and be careful not exceed recommended dosages.

Possibly Effective

High Fiber Diets. One oldest and most common methods of suppressing the appetite is to eat a high fiber diet. Ingesting high fiber

foods (fruits, vegetables) or fiber supplements increase the feeling of fullness (satiety). They typically allow you to feel full while ingesting fewer calories. Theoretically, maintaining a high fiber diet may serve to help decrease the amount of food you eat. In addition, high fiber diets/supplements have also been purported to help lower cholesterol and blood pressure as well as help diabetics manage glucose and insulin levels. Some of the research conducted on high fiber diets indicates that they provide some benefit, particularly in diabetic populations. For example, Raben et al²¹² reported that subjects maintaining a low fat/high fiber diet for 11 weeks lost about 3 lbs of weight and 3.5 lbs of fat. Other studies report either no significant effects or modest amounts of fat loss. High fiber/low fat diets have also been found to help reduce cholesterol. Consequently, although maintaining a low fat / high fiber diet may have some health benefits, they do not appear to promote a significant amount of weight or fat loss.

Calcium. Research has indicated that calcium modulates 1,25-dihydroxyvitamin D which serves to regulate intracellular calcium levels in fat cells²¹³⁻²¹⁵. Increasing dietary availability of calcium reduces 1,25-dihydroxyvitamin D and promotes reductions in fat mass in animals²¹³⁻²¹⁵. Dietary calcium has been shown to suppress fat metabolism and weight gain during periods of high caloric intake^{213, 216}. Further, increasing calcium intake has been shown to increase fat metabolism and preserve thermogenesis during caloric restriction^{213, 216, 217}. In support of this theory, Davies and colleagues²¹⁸ reported that dietary calcium was negatively correlated to weight and that calcium supplementation (1,000 mg/d) accounted for an 8 kg weight loss over a 4 yr period. Additionally, Zemel and associates²¹³ reported that supplemental calcium (800 mg/d) or high dietary intake of calcium (1,200 – 1,300 mg/d) during a 24-week weight loss program promoted significantly greater weight loss (26-70%) and dual energy x-ray absorptiometer (DEXA) determined fat mass

loss (38-64%) compared to subjects on a low calcium diet (400-500 mg/d). These findings and others suggest a strong relationship between calcium intake and fat loss.

Phosphates. The role of sodium and calcium phosphate on energy metabolism and exercise performance has been studied for decades³⁶. These studies have revealed that sodium phosphate supplementation appears to possess ergogenic properties particularly in endurance exercise events^{219, 220}. More recently, phosphate supplementation has also been suggested to affect energy expenditure. For example, Kaciuba-Uscilko and colleagues²²¹ reported that phosphate supplementation during a 4-week weight loss program increased resting metabolic rate (RMR) and respiratory exchange ratio (suggesting greater carbohydrate utilization and caloric expenditure) during submaximal cycling exercise. In addition, Nazar and coworkers²²² reported that phosphate supplementation during an 8-week weight loss program increased RMR by 12-19% and prevented a normal decline in thyroid hormones. Although the rate of weight loss was similar in this trial, results suggest that phosphate supplementation may influence metabolic rate possibly by affecting thyroid hormones. Consequently, it is possible that phosphate could serve as a potential thermogenic nutrient in non-ephedrine based supplements. Additional research is necessary to test this hypothesis.

Green Tea Extract. Green tea is one of the more interesting herbal supplements that has recently been suggested to affect weight loss. Green tea contains high amounts of caffeine and catechin polyphenols. Research suggests that catechin polyphenols possess antioxidant properties²²³. In addition, green tea has also been theorized to increase energy expenditure by stimulating brown adipose tissue thermogenesis. In support of this theory, Dulloo et al^{224, 225} reported that green tea supplementation in combination with caffeine (e.g., 50 mg caffeine and 90 mg epigallocatechin gallate taken 3-times per

day) significantly increased 24-hour energy expenditure and fat utilization in humans. The thermogenic effects of green tea supplementation were much greater than when an equivalent amount of caffeine was evaluated suggesting a synergistic effect. Theoretically, increases in energy expenditure may help individuals lose weight and/or manage body composition.

Calcium Pyruvate. Calcium Pyruvate is a supplement that hit the scene about five or six years ago with great promise. The theoretical rationale was based on studies from the early 1990s that reported that calcium pyruvate supplementation (16 – 25 g/d with or without dihydroxyacetone phosphate [DHAP]) promoted fat loss in overweight/obese patients following a medically supervised weight loss program²²⁶⁻²²⁸. Although the mechanism for these findings was unclear, the researchers speculated that it might be related to appetite suppression and/or altered carbohydrate and fat metabolism. Since calcium pyruvate is very expensive, several studies have attempted to determine whether ingesting smaller amounts of calcium pyruvate (6-10 g/d) affect body composition in untrained and trained populations. Results of these studies are mixed. Kalman and colleagues²²⁹ reported that calcium pyruvate supplementation (6 g/d for 6-weeks) significantly decreased body weight (-1.2 kg), body fat (-2.5 kg), and percent body fat (-2.7%). However, Stone and colleagues²³⁰ reported that pyruvate supplementation did not affect hydrostatically determined body composition during 5-weeks of in-season college football training. These findings indicate that although there is some supportive data indicating that calcium pyruvate supplementation may enhance fat loss when taken at high doses (6-16 g/d), there is no evidence that ingesting the doses typically found in pyruvate supplements (0.5 – 2 g/d) has any affect on body composition.

Too Early to Tell

Gymnema Sylvestre. *Gymnema Sylvestre* is a

relatively new supplement. It is purported to affect glucose and fat metabolism as well as inhibit sweet cravings. In support of these contentions, some recent data have been published by Shigematsu and colleagues^{231, 232} indicating that short and long-term oral supplementation of gymnema sylvestre in rats fed normal and high-fat diets may have some positive effects on fat metabolism, blood lipid levels, and/or weight gain/fat deposition. Although these findings are interesting, we are aware of no published studies that have evaluated the effects of gymnema sylvestre supplementation on lipid metabolism or body composition in humans. Consequently, more research is needed before conclusions can be drawn.

Chitosan. Chitosan has been marketed as a weight loss supplement for several years. It is purported to inhibit fat absorption and lower cholesterol. Several animal studies report decreased fat absorption, increased fecal fat content, and/or lower cholesterol following chitosan feedings²³³⁻²³⁶. However, the effects in humans appear to be less impressive. For example, although there is some data suggesting that chitosan supplementation may lower blood lipids in humans,²³⁷ other studies report no effects on fecal fat content²³⁸ or body composition alterations^{239, 240} when administered to people following their normal diet. It seems that people may be prone to eat more when they know they are taking a fat blocking supplement in a similar way people tend to eat more when they consume low-fat foods. Whether chitosan may promote greater amounts of fat loss when people are put on a controlled diet is unclear.

Non-Ephedra Containing Thermogenics. Since the safety of ephedra supplements has come into question, a number of supplement companies have been looking for alternatives to ephedra such as Citrus Aurantium or Bitter Orange (synephrine), thyroid stimulators, and various herbs and peppers (cayenne, black pepper, ginger root, etc)¹⁹¹. Of these, Citrus Aurantium (synephrine) appears to have the most promise^{241, 242}. Some studies suggest

that synephrine may increase metabolism without significantly affecting heart rate and blood pressure. However, it is unclear whether dietary supplementation of Citrus Aurantium may enhance weight loss. A number of thyroid stimulating supplements have also been marketed. Most contain nutrients (e.g., guggulsterones, 3, 5-Diiodo-L-Thyronine, etc.) believed to enhance the conversion of triiodothyronine (T3) to thyroxine (T4) or increase availability of T2 (diiodothyronine) or T3 which would theoretically increase basal metabolic rate (resting caloric expenditure) and promote weight loss^{243, 244}. However, while thyroid medications can effectively increase metabolic rate²⁴⁵, it is unclear whether these supplements can promote weight loss. Additionally, several of these types of supplements have been recently pulled by the FDA due to adverse health outcomes reported among people using these types of supplements particularly if they also contain usnic acid.

Phosphatidyl Choline (Lecithin). Choline is considered an essential nutrient that is needed for cell membrane integrity and to facilitate the movement of fats in and out of cells. It is also a component of the neurotransmitter acetylcholine and is needed for normal brain functioning, particularly in infants. For this reason, phosphatidyl choline (PC) has been purported as a potentially effective supplement to promote fat loss as well as improve neuromuscular function. There is some data from animal studies that supports the potential value of PC as a weight loss supplement²⁴⁶. There has also been some interest in determining the potential ergogenic value of choline supplementation during endurance exercise^{247, 248}. However, it is currently unclear whether PC supplementation affects body composition in humans.

Betaine. Betaine is a compound that is involved in the metabolism of choline and homocysteine. A number of studies have evaluated the effects of betaine feedings on

liver metabolism, fat metabolism, and fat deposition in animals^{249, 250}. There has also been interest in determining whether betaine supplementation may help lower homocysteine levels which has recently been identified as a marker of risk to heart disease²⁵¹. For this reason, betaine supplements have been marketed as a supplement designed to promote heart health as well as a weight loss. Although the potential theoretical rationale of betaine supplementation is interesting, it is currently unclear whether betaine supplementation may serve as an effective weight loss supplement in humans.

Coleus Forskohlii (Forskolin). Forskolin is another relatively new weight loss supplement. Forskolin is a plant native to India that has been used for centuries in traditional Ayurvedic medicine primarily to treat skin disorders and respiratory problems^{252, 253}. A considerable amount of research has evaluated the physiological and potential medical applications of forskolin over the last 25 years. Forskolin has been reported to reduce blood pressure, increase the hearts ability to contract, help inhibit platelet aggregation, improve lung function, and aid in the treatment of glaucoma²⁵²⁻²⁵⁴. With regard to weight loss, forskolin has been reported to increase cyclic AMP and thereby stimulate fat metabolism²⁵⁵⁻²⁵⁷.

Theoretically, forskolin may therefore serve as an effective weight loss supplement. In support of this theory, Sabinsa Corporation (the principle source for Forskolin in the U.S.) reported that forskolin supplementation (250 mg of a 10% forskolin extract taken twice daily for 8-weeks) administered in an open label manner to six overweight females promoted a 7.25 lbs loss in body weight and a 7.7% decrease bioelectrical impedance (BIA) determined body fat²⁵⁸. Although this was not a placebo controlled double blind study and BIA is not the most accurate method of assessing body composition, these preliminary findings provide some support to contentions that forskolin supplementation may promote fat loss. Another recent study suggested that supplementing the diet with

coelus forskohlii in overweight women helped maintain weight and was not associated with any clinically significant adverse events²⁵⁹. Additional research is needed before conclusions can be drawn.

Dehydroepiandrosterone (DHEA) and 7-Keto DHEA. Dehydroepiandrosterone (DHEA) and its sulfated conjugate DHEAS represent the most abundant adrenal steroids in circulation²⁶⁰. Although, DHEA is considered a weak androgen, it can be converted to the more potent androgens testosterone and dihydrotestosterone in tissues. In addition, DHEAS can be converted into androstenedione and testosterone. DHEA levels have been reported to decline with age in humans²⁶¹. The decline in DHEA levels with aging has been associated with increased fat accumulation and risk to heart disease²⁶². Since DHEA is a naturally occurring compound, it has been suggested that dietary supplementation of DHEA may help maintain DHEA availability, maintain and/or increase testosterone levels, reduce body fat accumulation, and/or reduce risk to heart disease as one ages^{260, 262}. Although animal studies have generally supported this theory, the effects of DHEA supplementation on body composition in human trials have been mixed. For example, Nestler and coworkers²⁶³ reported that DHEA supplementation (1,600 mg/d for 28-d) in untrained healthy males promoted a 31% reduction in percentage of body fat. However, Vogiatzi and associates²⁶⁴ reported that DHEA supplementation (40 mg/d for 8 wks) had no effect on body weight, percent body fat, or serum lipid levels in obese adolescents. More recently, 7-keto DHEA has been marketed as a potentially more effective form of DHEA. 7-keto DHEA is a precursor to DHEA that is believed to possess lipolytic properties. Although data are limited, Kalman and colleagues and coworkers²⁶⁵ reported that 7-keto DHEA supplementation (200 mg/d) during 8-weeks of training promoted a greater loss in body mass and fat mass while increasing T3. No significant effects were

observed on thyroid stimulating hormone (TSH), T4, or other hormones. Although more research is needed, these findings provide some support to contentions that 7-keto DHEA may serve as an effective weight loss supplement. However, additional research is needed before definitive conclusions can be made.

Psychotropic Nutrients/Herbs. This is a relatively new type of weight loss supplement category. Psychotropic nutrients/herbs often contain things like St. John's Wart, Kava, Ginkgo Biloba, Ginseng, and L-Tyrosine. They are believed to serve as naturally occurring antidepressants, relaxants, and mental stimulants. The theoretical rationale regarding weight loss is that they may help people fight depression or maintain mental alertness while dieting. Although a number of studies support potential role as naturally occurring psychotropics or stimulants, the potential value in promoting weight loss is unclear.

Apparently Ineffective

Chromium. Interest in chromium as a potential body composition modifier emanated from studies suggesting that chromium may enhance insulin sensitivity/glucose disposal in diabetics. Initial studies reported that chromium supplementation during resistance training improved fat loss and gains in lean body mass¹²⁸⁻¹³⁰. However, recent studies using more accurate methods of assessing body composition have mostly reported no effects on body composition in healthy non-diabetic individuals¹³¹⁻¹³⁷. For example, Walker and colleagues¹³² reported that chromium supplementation (200 µg/d for 14-weeks) did not affect body composition alterations during training in healthy wrestlers. Likewise, Lukaski et al¹³⁶ reported that 8-weeks of chromium supplementation during resistance training did not affect strength or DEXA determined body composition changes. Therefore, chromium supplementation does not appear to promote fat loss.

Conjugated Linoleic Acids (CLA). CLA is a term used to describe a group of positional and geometric isomers of linoleic acid that contain conjugated double bonds. Adding CLA to the diet has been reported to possess significant health benefits in animals^{138, 266}. In terms of weight loss, CLA feedings to animals have been reported to markedly decrease body fat accumulation^{138, 139, 143}. Consequently, CLA has been marketed as a health and weight loss supplement since the mid 1990s. Although basic research in animals is very promising, the effect of CLA supplementation in humans is less clear. There are some data suggesting that CLA supplementation may modestly promote fat loss and/or increases in lean mass^{144, 267-271}. However, other studies indicate that CLA supplementation (1.7 to 12 g/d for 4-weeks to 6-months) has limited to no effects on body composition alterations in untrained or trained populations^{144, 146, 147, 267, 272-274}. The reason for the discrepancy in research findings has been suggested to be due to differences in purity and the specific isomer studied. For instance, early studies in humans showing no effect used CLA that contained all 24 isomers. Today, most labs studying CLA use 50-50 mixtures containing the trans-10,cis-12 and cis-9,trans-11 isomers, the former of which being recently implicated in positively altering body composition. In our view, although CLA supplementation may have promise to promote general health, additional research is needed to determine if specific isomers of CLA may affect body composition in humans before conclusions can be made.

Garcinia Cambogia (HCA). HCA is a nutrient that has been hypothesized to increase fat oxidation by inhibiting citrate lyase and lipogenesis²⁷⁵. Theoretically, this may lead to greater fat burning and weight loss over time. Although there is some evidence that HCA may increase fat metabolism in animal studies, there is little to no evidence showing that HCA supplementation affects body composition in humans. For example, Ishihara et al²⁷⁶

reported that HCA supplementation spared carbohydrate utilization and promoted lipid oxidation during exercise in mice. However, Kriketos and associates²⁷⁷ reported that HCA supplementation (3 g/d for 3-days) did not affect resting or post-exercise energy expenditure or markers of lipolysis in healthy men. Likewise, Heymsfield and coworkers²⁷⁸ reported that HCA supplementation (1.5 g/d for 12-weeks) while maintaining a low fat/high fiber diet did not promote greater weight or fat loss than subjects on placebo. Finally, Mattes and colleagues²⁷⁹ reported that HCA supplementation (2.4 g/d for 12-weeks) did not affect appetite, energy intake, or weight loss. These findings suggest that HCA supplementation does not appear to promote fat loss in humans.

L-Carnitine. Carnitine serves as an important transporter of fatty acids from the cytosol into the mitochondria of the cell. Theoretically, increasing cellular levels of carnitine would thereby enhance transport of fats into the mitochondria and fat metabolism. For this reason, L-carnitine has been one of the most common nutrients found in various weight loss supplements. Over the years, a number of studies have been conducted on the effects of L-carnitine supplementation on fat metabolism, exercise capacity and body composition. Although there is some data showing that L-carnitine supplementation may be beneficial for some patient populations, most well controlled studies indicate that L-carnitine supplementation does not affect muscle carnitine content, fat metabolism, and/or weight loss in overweight or trained subjects²⁸⁰. For example, Villani et al²⁸¹ reported that L-carnitine supplementation (2 g/d for 8-weeks) did not affect weight loss, body composition, or markers of fat metabolism in overweight women.

Herbal Diuretics. This is a new type of supplement recently marketed as a natural way to promote weight loss. There is limited evidence that taraxacum officinale, verbena officinalis, lithospermum officinale,

equisetum arvense, arctostaphylos uva-ursi, arctium lappa and silene saxifraga infusion may affect diuresis in animals^{282, 283}. Two studies presented at the 2001 American College of Sports Medicine meeting^{284, 285} indicated that although herbal diuretics promoted a small amount of dehydration (about 0.3% in one day), they were not nearly as effective as a common diuretic drug (about 3.1% dehydration in one day). Consequently, although more research is needed, the potential value of herbal diuretics as a weight loss supplement appears limited.

Performance Enhancement Supplements

A number of nutritional supplements have been proposed to enhance exercise performance. Some of these nutrients have been described above. Table 3 categorizes the proposed ergogenic nutrients into apparently safe and effective, possibly effective, too early to tell, and apparently ineffective. Weight gain supplements purported to increase muscle mass may also have ergogenic properties if they also promote increases in strength. Similarly, some sports may benefit from reductions in fat mass. Therefore, weight loss supplements that help athletes manage body weight and/or fat mass may also possess some ergogenic benefit. The following describes which supplements may or may not affect performance that were not previously described. Based on this analysis, Table 4 summarizes the general nutritional recommendations for athletes and which dietary supplements may help power and endurance athletes.

Apparently Effective

Water and Sports Drinks. Preventing dehydration during exercise is one of the keys of maintaining exercise performance (particularly in hot/humid environments). People engaged in intense exercise or work in the heat need to frequently ingest water or sports drinks (e.g., 1-2 cups every 10 – 15 minutes). The goal should be not to lose

more than 2% of body weight during exercise (e.g., 180 lbs x 0.02 = 3.6 lbs). Sports drinks contain salt and carbohydrate. Studies show that ingestion of sports drinks during exercise in hot/humid environments can help prevent dehydration and improve endurance exercise capacity⁴³. Consequently, frequent ingestion of water and/or sports drinks during exercise is one of the easiest and most effective ergogenic aids.

Carbohydrate. General nutritional needs were discussed earlier. However, one of the best ergogenic aids available for active people is carbohydrate. Athletes and active individuals should consume a diet high in carbohydrate (e.g., 55 – 65% of calories or 5-8 grams/kg/day) in order to maintain muscle and liver carbohydrate stores¹. Additionally, ingesting a small amount of carbohydrate and protein 30-60 minutes prior to exercise and use of sports drinks during exercise can increase carbohydrate availability and improve exercise performance. Finally, ingesting carbohydrate and protein immediately following exercise can enhance carbohydrate storage and protein synthesis¹.

Creatine. Earlier we indicated that creatine supplementation is one of the best supplements available to increase muscle mass and strength during training. However, creatine has also been reported to improve exercise capacity in a variety of events⁵⁴. This is particularly true when performing high intensity, intermittent exercise such as multiple sets of weight lifting, repeated sprints, and/or exercise involving sprinting and jogging (e.g., soccer)⁵⁴. Although studies evaluating the ergogenic value of creatine on endurance exercise performance are mixed, endurance athletes may also theoretically benefit in several ways. For example, increasing creatine stores prior to carbohydrate loading (i.e., increasing dietary carbohydrate intake before competition in an attempt to maximize carbohydrate stores) has been shown to improve the ability to store carbohydrate²⁸⁶⁻²⁸⁸. Further, coingesting creatine with carbohydrate has been shown to

optimize creatine and carbohydrate loading²⁸⁹. Most endurance athletes also perform interval training (sprint or speed work) in an attempt to improve anaerobic threshold. Since creatine has been reported to enhance interval sprint performance, creatine supplementation during training may improve training adaptations in endurance athletes^{290, 291}. Finally, many endurance athletes lose weight during their competitive season. Creatine supplementation during training may help people maintain weight.

Sodium Phosphate. We previously mentioned that sodium phosphate supplementation may increase resting energy expenditure and therefore could serve as a potential weight loss nutrient. However, most research on sodium phosphate has actually evaluated the potential ergogenic value. A number of studies indicated that sodium phosphate supplementation (e.g., 1 gram taken 4 times daily for 3-6 days) can increase maximal oxygen uptake (i.e., maximal aerobic capacity) and anaerobic threshold by 5-10%^{219, 220, 292, 293}. These findings suggest that sodium phosphate may be highly effective in improving endurance exercise capacity. Other forms of phosphate (i.e., calcium phosphate, potassium phosphate) do not appear to possess ergogenic value.

Sodium Bicarbonate (Baking Soda). During high intensity exercise, acid (H⁺) and carbon dioxide (CO₂) accumulate in the muscle and blood. One of the ways you get rid of the acidity and CO₂ is to buffer the acid and CO₂ with bicarbonate ions. The acid and CO₂ are then removed in the lungs. Bicarbonate loading (e.g., 0.3 grams per kg taken 60-90 minutes prior to exercise or 5 grams taken 2 times per day for 5-days) has been shown to be an effective way to buffer acidity during high intensity exercise lasting 1-3 minutes in duration²⁹⁴⁻²⁹⁷. This can improve exercise capacity in events like the 400 - 800 m run or 100 – 200 m swim²⁹⁸. Although bicarbonate loading can improve exercise, some people have difficulty with their stomach tolerating

bicarbonate as it may cause gastrointestinal distress.

Caffeine. Caffeine is a naturally derived stimulant found in many nutritional supplements typically as Gaurana, Bissey Nut, or Kola. Caffeine can also be found in coffee, tea, soft drinks, energy drinks, and chocolate. Studies indicate that ingestion of caffeine (e.g., 3-9 mg/kg taken 30 – 90 minutes before exercise) can spare carbohydrate use during exercise and thereby improve endurance exercise capacity^{295, 299}. People who drink caffeinated drinks regularly, however, appear to experience less ergogenic benefits from caffeine³⁰⁰. Additionally, some concern has been expressed that ingestion of caffeine prior to exercise may contribute to dehydration although recent studies have not supported this concern³⁰¹⁻³⁰³. Caffeine doses above 9 mg/kg can result in urinary caffeine levels that surpass the doping threshold for many sport organizations. Suggestions that there is no ergogenic value to caffeine supplementation is not supported by the preponderance of available scientific studies.

Possibly Effective

Post-Exercise Carbohydrate and Protein. Ingesting carbohydrate and protein following exercise enhances carbohydrate storage and protein synthesis. Theoretically, ingesting carbohydrate and protein following exercise may lead to greater training adaptations. In support of this theory, Esmarck and coworkers⁹³ found that ingesting carbohydrate and protein immediately following exercise doubled training adaptations in comparison to waiting until 2-hours to ingest carbohydrate and protein. Additionally, Tarnopolsky and associates³⁰⁴ reported that post-exercise ingestion of carbohydrate with protein promoted as much strength gains as ingesting creatine with carbohydrate during training. These findings underscore the importance of post-exercise carbohydrate and protein ingestion.

Glutamine. As described above, glutamine has been shown to influence protein synthesis and help maintain the immune system. Theoretically, glutamine supplementation during training should enhance gains in strength and muscle mass as well as help athletes tolerate training to a better degree. Although there is some evidence that glutamine supplementation with protein can improve training adaptations, more research is needed to determine the ergogenic value in athletes.

Essential Amino Acids (EAA). Ingestion of 3-6 grams of EAA following resistance exercise has been shown to increase protein synthesis⁸⁴⁻⁹². Theoretically, ingestion of EAA after exercise should enhance gains in strength and muscle mass during training. While there is sound theoretical rationale, it is currently unclear whether following this strategy would lead to greater training adaptations and/or whether EAA supplementation would be better than simply ingesting carbohydrate and a quality protein following exercise.

Branched Chain Amino Acids (BCAA). Ingestion of BCAA (e.g., 6-10 grams per hour) with sports drinks during prolonged exercise would theoretically improve psychological perception of fatigue (i.e., central fatigue). Although there is strong rationale, the effects of BCAA supplementation on exercise performance is mixed with some studies suggesting an improvement and others showing no effect³⁶. More research is needed before conclusions can be drawn.

Calcium β -HMB. HMB supplementation has been reported to improve training adaptations in untrained individuals initiating training as well as help reduce muscle breakdown in runners. Theoretically, this should enhance training adaptations in athletes. However, most studies show little benefit of HMB supplementation in athletes.

Glycerol. Ingesting glycerol with water has been reported to increase fluid retention³⁰⁵. Theoretically, this should help athletes prevent dehydration during prolonged exercise and improve performance particularly if they are susceptible to dehydration. Although studies indicate that glycerol can significantly enhance body fluid, studies are mixed on whether it can improve exercise capacity^{52, 306-311}.

Ephedrine/Caffeine. Most research has evaluated the effects of ingesting ephedrine and caffeine (EC) supplements on weight loss. However, since ephedra and caffeine are stimulants and caffeine has been shown to have ergogenic properties, there has also been interest in the potential ergogenic value of EC. Recent research has shown that ingestion of low to moderate amounts of synthetic EC supplements generally improves endurance and high intensity exercise performance with no apparent adverse effects³¹²⁻³¹⁶. However, it is unclear whether dietary supplements containing botanical ephedrine (i.e., ephedra) and caffeine (e.g., kola nut) have similar effects on performance. Further, since most sport organizations ban use of ephedrine the potential use in athletes appears limited.

Too Early to Tell

A number of supplements purported to enhance performance and/or training adaptation fall under this category. This includes the weight gain and weight loss supplements listed in Table 3 as well as the following supplements not previously described in this category.

Medium Chain Triglycerides (MCT).

MCT's are shorter chain fatty acids that can easily enter the mitochondria of the cell and be converted to energy through fat metabolism³¹⁷. Studies are mixed as to whether MCT's can serve as an effective source of fat during exercise metabolism and/or improve exercise performance³¹⁸⁻³²².

Ribose. Ribose is a 3-carbon carbohydrate that is involved in the synthesis of adenosine triphosphate (ATP) in the muscle (the useable form of energy). Clinical studies have shown that ribose supplementation can increase exercise capacity in heart patients³²³⁻³²⁷. For this reason, ribose has been suggested to be an ergogenic aid for athletes. Although more research is needed, most studies show no ergogenic value of ribose supplementation on exercise capacity in health untrained or trained populations³²⁸⁻³³⁰.

Apparently Ineffective

Inosine. Inosine is a building block for DNA and RNA that is found in muscle. Inosine has a number of potentially important roles that may enhance training and/or exercise performance³³¹. Although there is some theoretical rationale, available studies indicate that inosine supplementation has no apparent affect on exercise performance capacity³³²⁻³³⁴.

Supplements to Promote General Health

In addition to the supplements previously described, several nutrients have been suggested to help athletes stay healthy during intense training. For example, the American Medical Association recently recommended that all Americans ingest a daily low-dose multivitamin in order to ensure that people get a sufficient amount of vitamins and minerals in their diet. Although one-a-day vitamin supplementation has not been found to improve exercise capacity in athletes, it may make sense to take a daily vitamin supplement for health reasons. Glucosamine and chondroitin have been reported to slow cartilage degeneration and reduce the degree of joint pain in active individuals which may help athletes postpone and/or prevent joint problems^{335, 336}. Vitamin C, glutamine, Echinacea, and zinc have been reported to enhance immune function^{42, 337-339}. Consequently, some sport nutritionists recommend that athletes who feel a cold coming on take these nutrients in order to

enhance immune function^{42, 337-339}. Similarly, nutrients such as vitamins E and C may help restore overwhelmed anti-oxidant defenses exhibited by athletes and reduce the risk of numerous chronic diseases³⁴⁰. Creatine, calcium β-HMB, BCAA, and L-carnitine have been shown to help athletes tolerate heavy training periods^{36, 74, 80, 81, 83, 341-344}. Finally, omega-3 fatty acids, in supplemental form, are now endorsed by the American Heart Association for heart health in certain individuals³⁴⁵. This supportive supplement position stems from: 1.) an inability to consume cardio-protective amounts by diet alone; and, 2.) the mercury contamination sometimes present in whole-food sources of DHA (docosahexaenoic acid) and EPA (eicosapentaenoic acid) found in fatty fish. Consequently, prudent use of these types of nutrients at various times during training may help athletes stay healthy and/or tolerate training to a greater degree³⁷.

SUMMARY

Numerous nutritional and herbal products are marketed to promote weight gain, weight loss, and/or improve performance. Most have a theoretical basis for use but little data supporting safety and efficacy in athletes. A number are heavily marketed despite data indicating that they do not affect body composition, performance, and/or training adaptations at the dosages recommended. It is in these particular situations that unsupported claims explicitly or implicitly endorsed by sport nutrition specialists constitute fraud and/ or “quackery”. Prudent

training, maintaining an energy balance and nutrient dense diet, proper timing of nutrient intake, and obtaining adequate rest are the cornerstones to enhancing performance and/or training adaptations. Use of a limited number of nutritional supplements that research has supported can help improve energy availability (e.g., sports drinks, carbohydrate, creatine, caffeine, etc) and/or promote recovery (carbohydrate, protein, essential amino acids, etc) can provide additional benefit in certain instances. The sport nutrition specialist should stay up to date regarding the role of nutrition on exercise so they can provide honest and accurate information to their students, clients, and/or athletes about the role of nutrition and dietary supplements on performance and training. Furthermore, the sport nutrition specialist should actively participate in exercise nutrition research; write unbiased scholarly reviews for journals and lay publications; help disseminate the latest research findings to the public so they can make informed decisions about appropriate methods of exercise, dieting, and/or whether various nutritional supplements can affect health, performance, and/or training; and, disclose any commercial or financial conflicts of interest during such promulgations to the public. Finally, sport nutrition specialists can challenge companies who sell exercise equipment and/or nutritional supplements to develop scientifically based products, conduct research on their products, and honestly market the results of studies so consumers can make informed decisions.

REFERENCES

1. Leutholtz B, Kreider RB. Exercise and Sport Nutrition. In: Wilson T, Temple N, eds. Nutritional Health. Totowa, NJ: Humana Press, Inc.; 2001:207 – 39.
2. Williams MH. Nutrition for Health, Fitness, and Sport. Dubuque, IA: ACB/McGraw-Hill; 1999.
3. FDA. Dietary Supplements. 2003: <http://www.cfsan.fda.gov/~dms/ds-faq.html>.
4. US RDA Recommendations.; Available: <http://www.lifestyler.com/jr/rdachart.htm>.
5. Beers MH, Berkow R. The Merck Manual. 17 ed: Merck Research Laboratories; 1999.
6. PDR for Nutritional Supplements. Montvale, NJ.: Medical Economics Co.; 2001.
7. PDR for Herbal Medicines. 2 ed. Montvale, NJ.: Medical Economics Co. Available: <http://physician.pdr.net/physician/static.htm?path=controlled/searchpdrherbal.htm>; 2000.
8. The Natural Health Encyclopedia. 2002; Available: <http://www.tnp.com/encyclopedia/>.

9. National Library of Medicine/Pub Med. 2002; Available: (<http://www.ncbi.nlm.nih.gov/PubMed/>).
10. Sherman WM, Jacobs KA, Leenders N. Carbohydrate metabolism during endurance exercise. In: Kreider RB, Fry AC, O'Toole ML, eds. *Overtraining in Sport*. Champaign: Human Kinetics Publishers; 1998:289-308.
11. Berning JR. Energy intake, diet, and muscle wasting. In: Kreider RB, Fry AC, O'Toole ML, eds. *Overtraining in Sport*. Champaign: Human Kinetics; 1998:275-88.
12. Kreider RB, Fry AC, O'Toole ML. *Overtraining in Sport*. Champaign, IL.: Human Kinetics Publishers; 1998.
13. Kreider RB. Physiological considerations of ultraendurance performance. *Int J Sport Nutr* 1991;1(1):3-27.
14. Brouns F, Saris WH, Beckers E, et al. Metabolic changes induced by sustained exhaustive cycling and diet manipulation. *Int J Sports Med* 1989;10 Suppl 1:S49-62.
15. Brouns F, Saris WH, Stroecken J, et al. Eating, drinking, and cycling. A controlled Tour de France simulation study, Part II. Effect of diet manipulation. *Int J Sports Med* 1989;10 Suppl 1:S41-8.
16. Lemon PW, Tarnopolsky MA, MacDougall JD, Atkinson SA. Protein requirements and muscle mass/strength changes during intensive training in novice bodybuilders. *J Appl Physiol* 1992;73(2):767-75.
17. Tarnopolsky MA, MacDougall JD, Atkinson SA. Influence of protein intake and training status on nitrogen balance and lean body mass. *J Appl Physiol* 1988;64(1):187-93.
18. Tarnopolsky MA, Atkinson SA, MacDougall JD, Chesley A, Phillips S, Schwarcz HP. Evaluation of protein requirements for trained strength athletes. *J Appl Physiol* 1992;73(5):1986-95.
19. Tarnopolsky MA. Protein and physical performance. *Curr Opin Clin Nutr Metab Care* 1999;2(6):533-7.
20. Kreider RB. Effects of protein and amino acid supplementation on athletic performance. *Sportscience* 1999. Available: [http://www.sportsci.org/jour/9901/rbk.html;3\(1\)](http://www.sportsci.org/jour/9901/rbk.html;3(1)).
21. Kreider RB, Kleiner SM. Protein supplements for athletes: need vs. convenience. *Your Patient & Fitness* 2000;14(6):12-8.
22. Bucci L, Unlu L. Proteins and amino acid supplements in exercise and sport. In: Driskell J, Wolinsky I, eds. *Energy-Yielding Macronutrients and Energy Metabolism in Sports Nutrition*. Boca Raton, FL: CRC Press; 2000:191-212.
23. Boirie Y, Dangin M, Gachon P, Vasson MP, Maubois JL, Beaufrere B. Slow and fast dietary proteins differently modulate postprandial protein accretion. *Proc Natl Acad Sci U S A* 1997;94(26):14930-5.
24. Boirie Y, Gachon P, Cordat N, Ritz P, Beaufrere B. Differential insulin sensitivities of glucose, amino acid, and albumin metabolism in elderly men and women. *J Clin Endocrinol Metab* 2001;86(2):638-44.
25. Boirie Y, Gachon P, Corny S, Fauquant J, Maubois JL, Beaufrere B. Acute postprandial changes in leucine metabolism as assessed with an intrinsically labeled milk protein. *Am J Physiol* 1996;271(6 Pt 1):E1083-91.
26. Venkatraman JT, Leddy J, Pendergast D. Dietary fats and immune status in athletes: clinical implications. *Med Sci Sports Exerc* 2000;32(7 Suppl):S389-95.
27. Dorgan JF, Judd JT, Longcope C, et al. Effects of dietary fat and fiber on plasma and urine androgens and estrogens in men: a controlled feeding study. *Am J Clin Nutr* 1996;64(6):850-5.
28. Hamalainen EK, Adlercreutz H, Puska P, Pietinen P. Decrease of serum total and free testosterone during a low-fat high-fibre diet. *J Steroid Biochem* 1983;18(3):369-70.
29. Reed MJ, Cheng RW, Simmonds M, Richmond W, James VH. Dietary lipids: an additional regulator of plasma levels of sex hormone binding globulin. *J Clin Endocrinol Metab* 1987;64(5):1083-5.
30. Fry AC, Kraemer WJ, Ramsey LT. Pituitary-adrenal-gonadal responses to high-intensity resistance exercise overtraining. *J Appl Physiol* 1998;85(6):2352-9.
31. Miller WC, Kocejka DM, Hamilton EJ. A meta-analysis of the past 25 years of weight loss research using diet, exercise or diet plus exercise intervention. *Int J Obes Relat Metab Disord* 1997;21(10):941-7.
32. Miller WC. Effective diet and exercise treatments for overweight and recommendations for intervention. *Sports Med* 2001;31(10):717-24.
33. Pirozzo S, Summerbell C, Cameron C, Glasziou P. Should we recommend low-fat diets for obesity? *Obes Rev* 2003;4(2):83-90.
34. Hu FB, Manson JE, Willett WC. Types of dietary fat and risk of coronary heart disease: a critical review. *J Am Coll Nutr* 2001;20(1):5-19.
35. Vessby B. Dietary fat, fatty acid composition in plasma and the metabolic syndrome. *Curr Opin Lipidol* 2003;14(1):15-9.
36. Kreider RB. Dietary supplements and the promotion of muscle growth with resistance exercise. *Sports Med* 1999;27(2):97-110.
37. Kreider RB. Nutritional Considerations of Overtraining. In: Stout JR, Antonio J, eds. *Sport Supplements: A Complete Guide to Physique and Athletic Enhancement*. Baltimore, MD: Lippincott, Williams & Wilkins; 2001:199-208.
38. Carli G, Bonifazi M, Lodi L, Lupo C, Martelli G, Viti A. Changes in the exercise-induced hormone response to branched chain amino acid administration. *Eur J Appl Physiol Occup Physiol* 1992;64(3):272-7.
39. Cade JR, Reese RH, Privette RM, Hommen NM, Rogers JL, Fregly MJ. Dietary intervention and training in swimmers. *Eur J Appl Physiol Occup Physiol* 1991;63(3-4):210-5.
40. Nieman DC, Fagoaga OR, Butterworth DE, et al. Carbohydrate supplementation affects blood granulocyte and monocyte trafficking but not function after 2.5 h of running. *Am J Clin Nutr* 1997;66(1):153-9.
41. Nieman DC. Influence of carbohydrate on the immune response to intensive, prolonged exercise. *Exerc Immunol Rev* 1998;4:64-76.
42. Nieman DC. Nutrition, exercise, and immune system function. *Clin Sports Med* 1999;18(3):537-48.
43. Burke LM. Nutritional needs for exercise in the heat. *Comp Biochem Physiol A Mol Integr Physiol* 2001;128(4):735-48.
44. Burke LM. Nutrition for post-exercise recovery. *Aust J Sci Med Sport* 1997;29(1):3-10.
45. Maughan RJ, Noakes TD. Fluid replacement and exercise stress. A brief review of studies on fluid replacement and some guidelines for the athlete. *Sports Med* 1991;12(1):16-31.
46. Zawadzki KM, Yaspelkis BB, 3rd, Ivy JL. Carbohydrate-protein complex increases the rate of muscle glycogen storage after exercise. *J Appl Physiol* 1992;72(5):1854-9.
47. Tarnopolsky MA, Bosman M, Macdonald JR, Vandeputte D, Martin J, Roy BD. Postexercise protein-carbohydrate and carbohydrate supplements increase muscle glycogen in men and women. *J Appl Physiol* 1997;83(6):1877-83.
48. Kraemer WJ, Volek JS, Bush JA, Putukian M, Sebastianelli WJ. Hormonal responses to consecutive days of heavy-resistance exercise with or without nutritional supplementation. *J Appl Physiol* 1998;85(4):1544-55.

49. Brouns F, Kovacs EM, Senden JM. The effect of different rehydration drinks on post-exercise electrolyte excretion in trained athletes. *Int J Sports Med* 1998;19(1):56-60.
50. Kovacs EM, Senden JM, Brouns F. Urine color, osmolality and specific electrical conductance are not accurate measures of hydration status during postexercise rehydration. *J Sports Med Phys Fitness* 1999;39(1):47-53.
51. Kovacs EM, Schmahl RM, Senden JM, Brouns F. Effect of high and low rates of fluid intake on post-exercise rehydration. *Int J Sport Nutr Exerc Metab* 2002;12(1):14-23.
52. Meyer LG, Horrigan DJ, Jr., Lotz WG. Effects of three hydration beverages on exercise performance during 60 hours of heat exposure. *Aviat Space Environ Med* 1995;66(11):1052-7.
53. Williams MH, Kreider R, Branch JD. *Creatine: The power supplement*. Champaign, IL: Human Kinetics Publishers; 1999.
54. Kreider RB. Effects of creatine supplementation on performance and training adaptations. *Mol Cell Biochem* 2003;244(1-2):89-94.
55. Volek JS, Duncan ND, Mazzetti SA, et al. Performance and muscle fiber adaptations to 12 weeks of creatine supplementation and heavy resistance training. *Medicine & Science in Sports & Exercise* 1999;31(5).
56. Willoughby DS, Rosene J. Effects of oral creatine and resistance training on myosin heavy chain expression. *Med Sci Sports Exerc* 2001;33(10):1674-81.
57. Willoughby DS, Rosene JM. Effects of oral creatine and resistance training on myogenic regulatory factor expression. *Med Sci Sports Exerc* 2003;35(6):923-9.
58. Kreider RB, Melton C, Rasmussen CJ, et al. Long-term creatine supplementation does not significantly affect clinical markers of health in athletes. *Mol Cell Biochem* 2003;244(1-2):95-104.
59. Graham AS, Hatton RC. Creatine: a review of efficacy and safety. *J Am Pharm Assoc (Wash)* 1999;39(6):803-10.
60. Juhn MS, Tarnopolsky M. Potential side effects of oral creatine supplementation: a critical review. *Clin J Sport Med* 1998;8(4):298-304.
61. Taes YE, Delanghe JR, Wuyts B, Van De Voorde J, Lameire NH. Creatine supplementation does not affect kidney function in an animal model with pre-existing renal failure. *Nephrol Dial Transplant* 2003;18(2):258-64.
62. Schilling BK, Stone MH, Utter A, et al. Creatine supplementation and health variables: a retrospective study. *Med Sci Sports Exerc* 2001;33(2):183-8.
63. Greenwood M, Kreider RB, Melton C, et al. Creatine supplementation during college football training does not increase the incidence of cramping or injury. *Mol Cell Biochem* 2003;244(1-2):83-8.
64. Greenwood M, Kreider R, Melton C, et al. Creatine supplementation during college football training does not increase the incidence of cramping or injury. *Mol Cell Biochem* 2002:In Press.
65. Watsford ML, Murphy AJ, Spinks WL, Walshe AD. Creatine supplementation and its effect on musculotendinous stiffness and performance. *J Strength Cond Res* 2003;17(1):26-33.
66. Nair KS, Matthews DE, Welle SL, Braiman T. Effect of leucine on amino acid and glucose metabolism in humans. *Metabolism* 1992;41(6):643-8.
67. Gallagher PM, Carrithers JA, Godard MP, Schulze KE, Trappe SW. Beta-hydroxy-beta-methylbutyrate ingestion, Part I: effects on strength and fat free mass. *Med Sci Sports Exerc* 2000;32(12):2109-15.
68. Gallagher PM, Carrithers JA, Godard MP, Schulze KE, Trappe SW. Beta-hydroxy-beta-methylbutyrate ingestion, part II: effects on hematology, hepatic and renal function. *Med Sci Sports Exerc* 2000;32(12):2116-9.
69. Nissen S, Sharp R, Ray M, et al. Effect of leucine metabolite beta-hydroxy-beta-methylbutyrate on muscle metabolism during resistance-exercise training. *J Appl Physiol* 1996;81(5):2095-104.
70. Pantan LB, Rathmacher JA, Baier S, Nissen S. Nutritional supplementation of the leucine metabolite beta-hydroxy-beta-methylbutyrate (hmb) during resistance training. *Nutrition* 2000;16(9):734-9.
71. Slater GJ, Jenkins D. Beta-hydroxy-beta-methylbutyrate (HMB) supplementation and the promotion of muscle growth and strength. *Sports Med* 2000;30(2):105-16.
72. Nissen S, Sharp RL, Pantan L, Vukovich M, Trappe S, Fuller JC, Jr. beta-hydroxy-beta-methylbutyrate (HMB) supplementation in humans is safe and may decrease cardiovascular risk factors. *J Nutr* 2000;130(8):1937-45.
73. Vukovich MD, Stubbs NB, Bohlken RM. Body composition in 70-year-old adults responds to dietary beta-hydroxy-beta-methylbutyrate similarly to that of young adults. *J Nutr* 2001;131(7):2049-52.
74. Knitter AE, Pantan L, Rathmacher JA, Petersen A, Sharp R. Effects of beta-hydroxy-beta-methylbutyrate on muscle damage after a prolonged run. *J Appl Physiol* 2000;89(4):1340-4.
75. Jowko E, Ostaszewski P, Jank M, et al. Creatine and beta-hydroxy-beta-methylbutyrate (HMB) additively increase lean body mass and muscle strength during a weight-training program. *Nutrition* 2001;17(7-8):558-66.
76. O'Connor DM, Crowe MJ. Effects of beta-hydroxy-beta-methylbutyrate and creatine monohydrate supplementation on the aerobic and anaerobic capacity of highly trained athletes. *J Sports Med Phys Fitness* 2003;43(1):64-8.
77. Kreider RB, Ferreira M, Wilson M, Almada AL. Effects of calcium beta-hydroxy-beta-methylbutyrate (HMB) supplementation during resistance-training on markers of catabolism, body composition and strength. *Int J Sports Med* 1999;20(8):503-9.
78. Slater G, Jenkins D, Logan P, et al. Beta-hydroxy-beta-methylbutyrate (HMB) supplementation does not affect changes in strength or body composition during resistance training in trained men. *Int J Sport Nutr Exerc Metab* 2001;11(3):384-96.
79. Ransone J, Neighbors K, Lefavi R, Chromiak J. The effect of beta-hydroxy beta-methylbutyrate on muscular strength and body composition in collegiate football players. *J Strength Cond Res* 2003;17(1):34-9.
80. Coombes JS, McNaughton LR. Effects of branched-chain amino acid supplementation on serum creatine kinase and lactate dehydrogenase after prolonged exercise. *J Sports Med Phys Fitness* 2000;40(3):240-6.
81. Schena F, Guerrini F, Tregnaghi P, Kayser B. Branched-chain amino acid supplementation during trekking at high altitude. The effects on loss of body mass, body composition, and muscle power. *Eur J Appl Physiol Occup Physiol* 1992;65(5):394-8.
82. Bigard AX, Lavier P, Ullmann L, Legrand H, Douce P, Guezennec CY. Branched-chain amino acid supplementation during repeated prolonged skiing exercises at altitude. *Int J Sport Nutr* 1996;6(3):295-306.
83. Candeloro N, Bertini I, Melchiorri G, De Lorenzo A. [Effects of prolonged administration of branched-chain amino acids on body composition and physical fitness]. *Minerva Endocrinol* 1995;20(4):217-23.
84. Tipton KD, Borsheim E, Wolf SE, Sanford AP, Wolfe RR. Acute response of net muscle protein balance reflects 24-h balance after exercise and amino acid ingestion. *Am J Physiol Endocrinol Metab* 2003;284(1):E76-89.

85. Wolfe RR. Regulation of muscle protein by amino acids. *J Nutr* 2002;132(10):3219S-24S.
86. Rasmussen BB, Tipton KD, Miller SL, Wolf SE, Wolfe RR. An oral essential amino acid-carbohydrate supplement enhances muscle protein anabolism after resistance exercise. *J Appl Physiol* 2000;88(2):386-92.
87. Tipton KD, Rasmussen BB, Miller SL, et al. Timing of amino acid-carbohydrate ingestion alters anabolic response of muscle to resistance exercise. *Am J Physiol Endocrinol Metab* 2001;281(2):E197-206.
88. Rasmussen BB, Wolfe RR, Volpi E. Oral and intravenously administered amino acids produce similar effects on muscle protein synthesis in the elderly. *J Nutr Health Aging* 2002;6(6):358-62.
89. Miller SL, Tipton KD, Chinkes DL, Wolf SE, Wolfe RR. Independent and combined effects of amino acids and glucose after resistance exercise. *Med Sci Sports Exerc* 2003;35(3):449-55.
90. Kobayashi H, Borsheim E, Anthony TG, et al. Reduced amino acid availability inhibits muscle protein synthesis and decreases activity of initiation factor eIF2B. *Am J Physiol Endocrinol Metab* 2003;284(3):E488-98.
91. Borsheim E, Tipton KD, Wolf SE, Wolfe RR. Essential amino acids and muscle protein recovery from resistance exercise. *Am J Physiol Endocrinol Metab* 2002;283(4):E648-57.
92. Biolo G, Williams BD, Fleming RY, Wolfe RR. Insulin action on muscle protein kinetics and amino acid transport during recovery after resistance exercise. *Diabetes* 1999;48(5):949-57.
93. Esmarck B, Andersen JL, Olsen S, Richter EA, Mizuno M, Kjaer M. Timing of postexercise protein intake is important for muscle hypertrophy with resistance training in elderly humans. *J Physiol* 2001;535(Pt 1):301-11.
94. Low SY, Taylor PM, Rennie MJ. Responses of glutamine transport in cultured rat skeletal muscle to osmotically induced changes in cell volume. *J Physiol* 1996;492 (Pt 3):877-85.
95. Rennie MJ, Khogali SE, Low SY, et al. Amino acid transport in heart and skeletal muscle and the functional consequences. *Biochem Soc Trans* 1996;24(3):869-73.
96. Rennie MJ, Ahmed A, Khogali SE, Low SY, Hundal HS, Taylor PM. Glutamine metabolism and transport in skeletal muscle and heart and their clinical relevance. *J Nutr* 1996;126(4 Suppl):1142S-9S.
97. Varnier M, Leese GP, Thompson J, Rennie MJ. Stimulatory effect of glutamine on glycogen accumulation in human skeletal muscle. *Am J Physiol* 1995;269(2 Pt 1):E309-15.
98. Antonio J, Street C. Glutamine: a potentially useful supplement for athletes. *Can J Appl Physiol* 1999;24(1):1-14.
99. Colker CM. Effects of supplemental protein on body composition and muscular strength in healthy athletic male adults. *Curr Ther Res* 2000;61(1):19-28.
100. Wernerman J, Hammarqvist F, Vinnars E. Alpha-ketoglutarate and postoperative muscle catabolism. *Lancet* 1990;335(8691):701-3.
101. Hammarqvist F, Wernerman J, von der Decken A, Vinnars E. Alanyl-glutamine counteracts the depletion of free glutamine and the postoperative decline in protein synthesis in skeletal muscle. *Ann Surg* 1990;212(5):637-44.
102. Antonio J, Stout JR. *Sport Supplements*. Philadelphia, PA: Lippincott, Williams and Wilkins.; 2001.
103. Mitch WE, Walser M, Sapir DG. Nitrogen sparing induced by leucine compared with that induced by its keto analogue, alpha-ketoisocaproate, in fasting obese man. *J Clin Invest* 1981;67(2):553-62.
104. Van Koeveering M, Nissen S. Oxidation of leucine and alpha-ketoisocaproate to beta-hydroxy-beta-methylbutyrate in vivo. *Am J Physiol* 1992;262(1 Pt 1):E27-31.
105. Slama K, Koudela K, Tenora J, Mathova A. Insect hormones in vertebrates: anabolic effects of 20-hydroxyecdysone in Japanese quail. *Experientia* 1996;52(7):702-6.
106. Slama K, Kodkova M. Insect hormones and bioanalogues: their effect on respiratory metabolism in *Dermestes vulpinus* L. (Coleoptera). *Biol Bull* 1975;148(2):320-32.
107. Tashmukhamedova MA, Almatov KT, Syrov VN, Sultanov MB, Abidov AA. [Effect of phytoecdysteroids and anabolic steroids on liver mitochondrial respiration and oxidative phosphorylation in alloxan diabetic rats]. *Nauchnye Doki Vyssh Shkoly Biol Nauki* 1985(9):37-9.
108. Syrov VN. [Mechanism of the anabolic action of phytoecdysteroids in mammals]. *Nauchnye Doki Vyssh Shkoly Biol Nauki* 1984(11):16-20.
109. Kholodova Y. Phytoecdysteroids: biological effects, application in agriculture and complementary medicine (as presented at the 14-th Ecdysone Workshop, July, 2000, Rapperswil, Switzerland). *Ukr Biokhim Zh* 2001;73(3):21-9.
110. Bowers CY. Growth hormone-releasing peptide (GHRP). *Cell Mol Life Sci* 1998;54(12):1316-29.
111. Camanni F, Ghigo E, Arvat E. Growth hormone-releasing peptides and their analogs. *Front Neuroendocrinol* 1998;19(1):47-72.
112. Messina M, Messina V. Soyfoods, soybean isoflavones, and bone health: a brief overview. *J Ren Nutr* 2000;10(2):63-8.
113. Messina M. Soyfoods and soybean phyto-oestrogens (isoflavones) as possible alternatives to hormone replacement therapy (HRT). *Eur J Cancer* 2000;36 Suppl 4:S71-2.
114. de Aloysio D, Gambacciani M, Altieri P, et al. Bone density changes in postmenopausal women with the administration of isoflavone alone or in association with low-dose ERT. *Gynecol Endocrinol* 1997;11(4):289-93.
115. US Patent 3949085: Anabolic-weight-gain promoting compositions containing isoflavone derivatives and method using same. Available at <http://www.delphion.com/details?pn=US03949085>.
116. US Patent 4163746: Metabolic 5-methyl-isoflavone-derivatives, process for the preparation thereof and compositions containing the same. Available at: <http://www.delphion.com/details?pn=US04163746>.
117. Chetlin RD, Yeater RA, Ullrich IH, Hornsby WG, Malanga CJ, Byrner RW. The effect of ornithine alpha-ketoglutarate (OKG) on healthy, weight trained men. *J Exerc Physiol Online* 2000;3(4):Available: www.css.edu/users/tboone2/asep/ChetlinV.pdf.
118. Gonzalez-Cadavid NF, Taylor WE, Yarasheski K, et al. Organization of the human myostatin gene and expression in healthy men and HIV-infected men with muscle wasting. *Proc Natl Acad Sci U S A* 1998;95(25):14938-43.
119. McPherron AC, Lee SJ. Double muscling in cattle due to mutations in the myostatin gene. *Proc Natl Acad Sci U S A* 1997;94(23):12457-61.
120. McPherron AC, Lawler AM, Lee SJ. Regulation of skeletal muscle mass in mice by a new TGF-beta superfamily member. *Nature* 1997;387(6628):83-90.
121. Grobet L, Martin LJ, Poncelet D, et al. A deletion in the bovine myostatin gene causes the double-muscling phenotype in cattle. *Nat Genet* 1997;17(1):71-4.

122. Kambadur R, Sharma M, Smith TP, Bass JJ. Mutations in myostatin (GDF8) in double-muscled Belgian Blue and Piedmontese cattle. *Genome Res* 1997;7(9):910-6.
123. Ivey FM, Roth SM, Ferrell RE, et al. Effects of age, gender, and myostatin genotype on the hypertrophic response to heavy resistance strength training. *J Gerontol A Biol Sci Med Sci* 2000;55(11):M641-8.
124. Carlson CJ, Booth FW, Gordon SE. Skeletal muscle myostatin mRNA expression is fiber-type specific and increases during hindlimb unloading. *Am J Physiol* 1999;277(2 Pt 2):R601-6.
125. Brilla LR, Conte V. Effects of a novel zing-magnesium formulation on hormones and strength. *J Exerc Physiol Online* 2000;3(4):Available: www.css.edu/users/tboone2/asep/BrillaV.pdf.
126. Green NR, Ferrando AA. Plasma boron and the effects of boron supplementation in males. *Environ Health Perspect* 1994;102 Suppl 7:73-7.
127. Ferrando AA, Green NR. The effect of boron supplementation on lean body mass, plasma testosterone levels, and strength in male bodybuilders. *Int J Sport Nutr* 1993;3(2):140-9.
128. Evans GW. The effect of chromium picolinate on insulin controlled parameters in humans. *Int Biosc Med Res* 1989;11:163-80.
129. Hasten DL, Rome EP, Franks BD, Hegsted M. Effects of chromium picolinate on beginning weight training students. *Int J Sport Nutr* 1992;2(4):343-50.
130. Grant KE, Chandler RM, Castle AL, Ivy JL. Chromium and exercise training: effect on obese women. *Med Sci Sports Exerc* 1997;29(8):992-8.
131. Campbell WW, Joseph LJ, Davey SL, Cyr-Campbell D, Anderson RA, Evans WJ. Effects of resistance training and chromium picolinate on body composition and skeletal muscle in older men. *J Appl Physiol* 1999;86(1):29-39.
132. Walker LS, Bemben MG, Bemben DA, Knehans AW. Chromium picolinate effects on body composition and muscular performance in wrestlers. *Med Sci Sports Exerc* 1998;30(12):1730-7.
133. Livolsi JM, Adams GM, Laguna PL. The effect of chromium picolinate on muscular strength and body composition in women athletes. *J Strength Cond Res* 2001;15(2):161-6.
134. Volpe SL, Huang HW, Larpadisorn K, Lesser, II. Effect of chromium supplementation and exercise on body composition, resting metabolic rate and selected biochemical parameters in moderately obese women following an exercise program. *J Am Coll Nutr* 2001;20(4):293-306.
135. Hallmark MA, Reynolds TH, DeSouza CA, Dotson CO, Anderson RA, Rogers MA. Effects of chromium and resistive training on muscle strength and body composition. *Med Sci Sports Exerc* 1996;28(1):139-44.
136. Lukaski HC, Bolonchuk WW, Siders WA, Milne DB. Chromium supplementation and resistance training: effects on body composition, strength, and trace element status of men. *Am J Clin Nutr* 1996;63(6):954-65.
137. Clancy SP, Clarkson PM, DeCheke ME, et al. Effects of chromium picolinate supplementation on body composition, strength, and urinary chromium loss in football players. *Int J Sport Nutr* 1994;4(2):142-53.
138. Pariza MW, Park Y, Cook ME. Conjugated linoleic acid and the control of cancer and obesity. *Toxicol Sci* 1999;52(2 Suppl):107-10.
139. Pariza MW, Park Y, Cook ME. Mechanisms of action of conjugated linoleic acid: evidence and speculation. *Proc Soc Exp Biol Med* 2000;223(1):8-13.
140. Pariza MW, Park Y, Cook ME. The biologically active isomers of conjugated linoleic acid. *Prog Lipid Res* 2001;40(4):283-98.
141. DeLany JP, Blohm F, Truett AA, Scimeca JA, West DB. Conjugated linoleic acid rapidly reduces body fat content in mice without affecting energy intake. *Am J Physiol* 1999;276(4 Pt 2):R1172-9.
142. DeLany JP, West DB. Changes in body composition with conjugated linoleic acid. *J Am Coll Nutr* 2000;19(4):487S-93S.
143. Park Y, Albright KJ, Liu W, Storkson JM, Cook ME, Pariza MW. Effect of conjugated linoleic acid on body composition in mice. *Lipids* 1997;32(8):853-8.
144. Blankson H, Stakkestad JA, Fagertun H, Thom E, Wadstein J, Gudmundsen O. Conjugated linoleic acid reduces body fat mass in overweight and obese humans. *J Nutr* 2000;130(12):2943-8.
145. Gaullier JM, Berven G, Blankson H, Gudmundsen O. Clinical trial results support a preference for using CLA preparations enriched with two isomers rather than four isomers in human studies. *Lipids* 2002;37(11):1019-25.
146. Zambell KL, Keim NL, Van Loan MD, et al. Conjugated linoleic acid supplementation in humans: effects on body composition and energy expenditure. *Lipids* 2000;35(7):777-82.
147. Kreider RB, Ferreira MP, Greenwood M, Wilson M, Almada AL. Effects of conjugated linoleic acid supplementation during resistance training on body composition, bone density, strength, and selected hematological markers. *J Strength Cond Res* 2002;16(3):325-34.
148. Wheeler KB, Garleb KA. Gamma oryzanol-plant sterol supplementation: metabolic, endocrine, and physiologic effects. *Int J Sport Nutr* 1991;1(2):170-7.
149. Fry AC, Bonner E, Lewis DL, Johnson RL, Stone MH, Kraemer WJ. The effects of gamma-oryzanol supplementation during resistance exercise training. *Int J Sport Nutr* 1997;7(4):318-29.
150. Yarasheski KE. Growth hormone effects on metabolism, body composition, muscle mass, and strength. *Exerc Sport Sci Rev* 1994;22:285-312.
151. Lukas SE. Current perspectives on anabolic-androgenic steroid abuse. *Trends Pharmacol Sci* 1993;14(2):61-8.
152. Wagner JC. Enhancement of athletic performance with drugs. An overview. *Sports Med* 1991;12(4):250-65.
153. Limbird TJ. Anabolic steroids in the training and treatment of athletes. *Compr Ther* 1985;11(1):25-30.
154. Kuhn CM. Anabolic steroids. *Recent Prog Horm Res* 2002;57:411-34.
155. Smart T. Other therapies for wasting. *GMHC Treat Issues* 1995;9(5):7-8, 12.
156. Casaburi R. Skeletal muscle dysfunction in chronic obstructive pulmonary disease. *Med Sci Sports Exerc* 2001;33(7 Suppl):S662-70.
157. Hayes VY, Urban RJ, Jiang J, Marcell TJ, Helgeson K, Mauras N. Recombinant human growth hormone and recombinant human insulin-like growth factor I diminish the catabolic effects of hypogonadism in man: metabolic and molecular effects. *J Clin Endocrinol Metab* 2001;86(5):2211-9.
158. Newshan G, Leon W. The use of anabolic agents in HIV disease. *Int J STD AIDS* 2001;12(3):141-4.
159. Tenover JS. Androgen replacement therapy to reverse and/or prevent age-associated sarcopenia in men. *Baillieres Clin*

- Endocrinol Metab 1998;12(3):419-25.
160. Bross R, Casaburi R, Storer TW, Bhasin S. Androgen effects on body composition and muscle function: implications for the use of androgens as anabolic agents in sarcopenic states. *Baillieres Clin Endocrinol Metab* 1998;12(3):365-78.
 161. Casaburi R. Rationale for anabolic therapy to facilitate rehabilitation in chronic obstructive pulmonary disease. *Baillieres Clin Endocrinol Metab* 1998;12(3):407-18.
 162. Johansen KL, Mulligan K, Schambelan M. Anabolic effects of nandrolone decanoate in patients receiving dialysis: a randomized controlled trial. *Jama* 1999;281(14):1275-81.
 163. Sattler FR, Jaque SV, Schroeder ET, et al. Effects of pharmacological doses of nandrolone decanoate and progressive resistance training in immunodeficient patients infected with human immunodeficiency virus. *J Clin Endocrinol Metab* 1999;84(4):1268-76.
 164. Beiner JM, Jokl P, Cholewicki J, Panjabi MM. The effect of anabolic steroids and corticosteroids on healing of muscle contusion injury. *Am J Sports Med* 1999;27(1):2-9.
 165. Ferreira IM, Verreschi IT, Nery LE, et al. The influence of 6 months of oral anabolic steroids on body mass and respiratory muscles in undernourished COPD patients. *Chest* 1998;114(1):19-28.
 166. Bhasin S, Storer TW, Berman N, et al. Testosterone replacement increases fat-free mass and muscle size in hypogonadal men. *J Clin Endocrinol Metab* 1997;82(2):407-13.
 167. Ferrando AA, Sheffield-Moore M, Paddon-Jones D, Wolfe RR, Urban RJ. Differential anabolic effects of testosterone and amino acid feeding in older men. *J Clin Endocrinol Metab* 2003;88(1):358-62.
 168. Meeuwse IB, Samson MM, Duursma SA, Verhaar HJ. Muscle strength and tibolone: a randomised, double-blind, placebo-controlled trial. *Bjog* 2002;109(1):77-84.
 169. King DS, Sharp RL, Vukovich MD, et al. Effect of oral androstenedione on serum testosterone and adaptations to resistance training in young men: a randomized controlled trial. *Jama* 1999;281(21):2020-8.
 170. Carter WJ. Effect of anabolic hormones and insulin-like growth factor-I on muscle mass and strength in elderly persons. *Clin Geriatr Med* 1995;11(4):735-48.
 171. Soe M, Jensen KL, Gluud C. [The effect of anabolic androgenic steroids on muscle strength, body weight and lean body mass in body-building men]. *Ugeskr Laeger* 1989;151(10):610-3.
 172. Griggs RC, Pandya S, Florence JM, et al. Randomized controlled trial of testosterone in myotonic dystrophy. *Neurology* 1989;39(2 Pt 1):219-22.
 173. Crist DM, Stackpole PJ, Peake GT. Effects of androgenic-anabolic steroids on neuromuscular power and body composition. *J Appl Physiol* 1983;54(2):366-70.
 174. Ward P. The effect of an anabolic steroid on strength and lean body mass. *Med Sci Sports* 1973;5(4):277-82.
 175. Varriale P, Mirzai-tehrane M, Sedighi A. Acute myocardial infarction associated with anabolic steroids in a young HIV-infected patient. *Pharmacotherapy* 1999;19(7):881-4.
 176. Kibble MW, Ross MB. Adverse effects of anabolic steroids in athletes. *Clin Pharm* 1987;6(9):686-92.
 177. Gruber AJ, Pope HG, Jr. Psychiatric and medical effects of anabolic-androgenic steroid use in women. *Psychother Psychosom* 2000;69(1):19-26.
 178. Lamb DR. Anabolic steroids in athletics: how well do they work and how dangerous are they? *Am J Sports Med* 1984;12(1):31-8.
 179. Salke RC, Rowland TW, Burke EJ. Left ventricular size and function in body builders using anabolic steroids. *Med Sci Sports Exerc* 1985;17(6):701-4.
 180. Broeder CE, Quindry J, Brittingham K, et al. The Andro Project: physiological and hormonal influences of androstenedione supplementation in men 35 to 65 years old participating in a high-intensity resistance training program. *Arch Intern Med* 2000;160(20):3093-104.
 181. Ballantyne CS, Phillips SM, MacDonald JR, Tarnopolsky MA, MacDougall JD. The acute effects of androstenedione supplementation in healthy young males. *Can J Appl Physiol* 2000;25(1):68-78.
 182. Brown GA, Vukovich MD, Sharp RL, Reifnath TA, Parsons KA, King DS. Effect of oral DHEA on serum testosterone and adaptations to resistance training in young men. *J Appl Physiol* 1999;87(6):2274-83.
 183. van Gammeren D, Falk D, Antonio J. Effects of norandrostenedione and norandrostenediol in resistance-trained men. *Nutrition* 2002;18(9):734-7.
 184. Brown GA, Martini ER, Roberts BS, Vukovich MD, King DS. Acute hormonal response to sublingual androstenediol intake in young men. *J Appl Physiol* 2002;92(1):142-6.
 185. Van Gammeren D, Falk D, Antonio J. The effects of supplementation with 19-nor-4-androstene-3,17-dione and 19-nor-4-androstene-3,17-diol on body composition and athletic performance in previously weight-trained male athletes. *Eur J Appl Physiol* 2001;84(5):426-31.
 186. Pipe A. Effects of testosterone precursor supplementation on intensive weight training. *Clin J Sport Med* 2001;11(2):126.
 187. Brown GA, Vukovich MD, Martini ER, et al. Effects of androstenedione-herbal supplementation on serum sex hormone concentrations in 30- to 59-year-old men. *Int J Vitam Nutr Res* 2001;71(5):293-301.
 188. Antonio J, Uelmen J, Rodriguez R, Earnest C. The effects of Tribulus terrestris on body composition and exercise performance in resistance-trained males. *Int J Sport Nutr Exerc Metab* 2000;10(2):208-15.
 189. Fawcett JP, Farquhar SJ, Walker RJ, Thou T, Lowe G, Goulding A. The effect of oral vanadyl sulfate on body composition and performance in weight-training athletes. *Int J Sport Nutr* 1996;6(4):382-90.
 190. Fawcett JP, Farquhar SJ, Thou T, Shand BI. Oral vanadyl sulphate does not affect blood cells, viscosity or biochemistry in humans. *Pharmacol Toxicol* 1997;80(4):202-6.
 191. Kreider RB. New weight-control options. *Functional Foods & Nutraceuticals* 2002;July/August:34-42.
 192. Hoie LH, Bruusgaard D, Thom E. Reduction of body mass and change in body composition on a very low calorie diet. *Int J Obes Relat Metab Disord* 1993;17(1):17-20.
 193. Bryner RW, Ullrich IH, Sauer J, et al. Effects of resistance vs. aerobic training combined with an 800 calorie liquid diet on lean body mass and resting metabolic rate. *J Am Coll Nutr* 1999;18(2):115-21.
 194. Kern PA, Trozzolino L, Wolfe G, Purdy L. Combined use of behavior modification and very low-calorie diet in weight loss and weight maintenance. *Am J Med Sci* 1994;307(5):325-8.
 195. Baba NH, Sawaya S, Torbay N, Habbal Z, Azar S, Hashim SA. High protein vs high carbohydrate hypoenergetic diet for the

- treatment of obese hyperinsulinemic subjects. *Int J Obes Relat Metab Disord* 1999;23(11):1202-6.
196. Skov AR, Toubro S, Ronn B, Holm L, Astrup A. Randomized trial on protein vs carbohydrate in ad libitum fat reduced diet for the treatment of obesity. *Int J Obes Relat Metab Disord* 1999;23(5):528-36.
 197. Toubro S, Astrup AV. [A randomized comparison of two weight-reducing diets. Calorie counting versus low-fat carbohydrate-rich ad libitum diet]. *Ugeskr Laeger* 1998;160(6):816-20.
 198. Aoyama T, Fukui K, Takamatsu K, Hashimoto Y, Yamamoto T. Soy protein isolate and its hydrolysate reduce body fat of dietary obese rats and genetically obese mice (yellow KK). *Nutrition* 2000;16(5):349-54.
 199. Reaven GM. Diet and Syndrome X. *Curr Atheroscler Rep* 2000;2(6):503-7.
 200. Boozer CN, Daly PA, Homel P, et al. Herbal ephedra/caffeine for weight loss: a 6-month randomized safety and efficacy trial. *Int J Obes Relat Metab Disord* 2002;26(5):593-604.
 201. Boozer CN, Nasser JA, Heymsfield SB, Wang V, Chen G, Solomon JL. An herbal supplement containing Ma Huang-Guarana for weight loss: a randomized, double-blind trial. *Int J Obes Relat Metab Disord* 2001;25(3):316-24.
 202. Boozer C, Daly P, Homel P, et al. Herbal ephedra/caffeine for weight loss: a 6-month randomized safety and efficacy trial. *Int J Obesity* 2002;26:593-604.
 203. Boozer C, Nasser J, SB H, Wang V, Chen G, Solomon J. An herbal supplement containing Ma Huang-Guarana for weight loss: a randomized, double-blind trial. *Int J Obes Relat Metab Disord* 2001;25:316-24.
 204. Molnar D, Torok K, Erhardt E, Jeges S. Safety and efficacy of treatment with an ephedrine/caffeine mixture. The first double-blind placebo-controlled pilot study in adolescents. *Int J Obes Relat Metab Disord* 2000;24(12):1573-8.
 205. Molnar D. Effects of ephedrine and aminophylline on resting energy expenditure in obese adolescents. *Int J Obes Relat Metab Disord* 1993;17 Suppl 1:S49-52.
 206. Greenway FL. The safety and efficacy of pharmaceutical and herbal caffeine and ephedrine use as a weight loss agent. *Obes Rev* 2001;2(3):199-211.
 207. Greenway FL, Raum WJ, DeLany JP. The effect of an herbal dietary supplement containing ephedrine and caffeine on oxygen consumption in humans. *J Altern Complement Med* 2000;6(6):553-5.
 208. Greenway FL, Ryan DH, Bray GA, Rood JC, Tucker EW, Smith SR. Pharmaceutical cost savings of treating obesity with weight loss medications. *Obes Res* 1999;7(6):523-31.
 209. Greenway F, Herber D, Raum W, Morales S. Double-blind, randomized, placebo-controlled clinical trials with non-prescription medications for the treatment of obesity. *Obes Res* 1999;7(4):370-8.
 210. Bent S, Tiedt T, Odden M, Shlipak M. The relative safety of ephedra compared with other herbal products. *Ann Intern Med* 2003; Available: <http://www.acponline.org/journals/annals/ephedra.htm>.
 211. Fleming GA. The FDA, regulation, and the risk of stroke. *N Engl J Med* 2000;343(25):1886-7.
 212. Raben A, Jensen ND, Marckmann P, Sandstrom B, Astrup AV. [Spontaneous weight loss in young subjects of normal weight after 11 weeks of unrestricted intake of a low-fat/high-fiber diet]. *Ugeskr Laeger* 1997;159(10):1448-53.
 213. Zemel M, Thompson W, Zemel P, et al. Dietary calcium and dairy products accelerate weight and fat-loss during energy restriction in obese adults. *Clin Nutri* 2002;75.
 214. Zemel MB. Role of dietary calcium and dairy products in modulating adiposity. *Lipids* 2003;38(2):139-46.
 215. Zemel MB. Mechanisms of dairy modulation of adiposity. *J Nutr* 2003;133(1):252S-6S.
 216. Zemel MB, Shi H, Greer B, Dirienzo D, Zemel PC. Regulation of adiposity by dietary calcium. *Faseb J* 2000;14(9):1132-8.
 217. Zemel MB. Regulation of adiposity and obesity risk by dietary calcium: mechanisms and implications. *J Am Coll Nutr* 2002;21(2):146S-51S.
 218. Davies KM, Heaney RP, Recker RR, et al. Calcium intake and body weight. *J Clin Endocrinol Metab* 2000;85(12):4635-8.
 219. Kreider RB, Miller GW, Schenck D, et al. Effects of phosphate loading on metabolic and myocardial responses to maximal and endurance exercise. *Int J Sport Nutr* 1992;2(1):20-47.
 220. Kreider RB, Miller GW, Williams MH, Somma CT, Nasser TA. Effects of phosphate loading on oxygen uptake, ventilatory anaerobic threshold, and run performance. *Med Sci Sports Exerc* 1990;22(2):250-6.
 221. Kaciuba-Uscilko H, Nazar K, Chwalbinska-Moneta J, et al. Effect of phosphate supplementation on metabolic and neuroendocrine responses to exercise and oral glucose load in obese women during weight reduction. *J Physiol Pharmacol* 1993;44(4):425-40.
 222. Nazar K, Kaciuba-Uscilko H, Szczepanik J, et al. Phosphate supplementation prevents a decrease of triiodothyronine and increases resting metabolic rate during low energy diet. *J Physiol Pharmacol* 1996;47(2):373-83.
 223. Nakagawa K, Ninomiya M, Okubo T, et al. Tea catechin supplementation increases antioxidant capacity and prevents phospholipid hydroperoxidation in plasma of humans. *J Agric Food Chem* 1999;47(10):3967-73.
 224. Dulloo AG, Seydoux J, Girardier L, Chantre P, Vandermander J. Green tea and thermogenesis: interactions between catechin-polyphenols, caffeine and sympathetic activity. *Int J Obes Relat Metab Disord* 2000;24(2):252-8.
 225. Dulloo AG, Duret C, Rohrer D, et al. Efficacy of a green tea extract rich in catechin polyphenols and caffeine in increasing 24-h energy expenditure and fat oxidation in humans. *Am J Clin Nutr* 1999;70(6):1040-5.
 226. Stanko RT, Arch JE. Inhibition of regain in body weight and fat with addition of 3-carbon compounds to the diet with hyperenergetic refeeding after weight reduction. *Int J Obes Relat Metab Disord* 1996;20(10):925-30.
 227. Stanko RT, Tietze DL, Arch JE. Body composition, energy utilization, and nitrogen metabolism with a severely restricted diet supplemented with dihydroxyacetone and pyruvate. *Am J Clin Nutr* 1992;55(4):771-6.
 228. Stanko RT, Reynolds HR, Hoyson R, Janosky JE, Wolf R. Pyruvate supplementation of a low-cholesterol, low-fat diet: effects on plasma lipid concentrations and body composition in hyperlipidemic patients. *Am J Clin Nutr* 1994;59(2):423-7.
 229. Kalman D, Colker CM, Wilets I, Roufs JB, Antonio J. The effects of pyruvate supplementation on body composition in overweight individuals. *Nutrition* 1999;15(5):337-40.
 230. Stone MH, Sanborn K, Smith LL, et al. Effects of in-season (5 weeks) creatine and pyruvate supplementation on anaerobic performance and body composition in American football players. *Int J Sport Nutr* 1999;9(2):146-65.
 231. Shigematsu N, Asano R, Shimosaka M, Okazaki M. Effect of administration with the extract of *Gymnema sylvestris* R. Br leaves on lipid metabolism in rats. *Biol Pharm Bull* 2001;24(6):713-7.
 232. Shigematsu N, Asano R, Shimosaka M, Okazaki M. Effect of long term-administration with *Gymnema sylvestris* R. BR on plasma and liver lipid in rats. *Biol Pharm Bull* 2001;24(6):643-9.
 233. Gallaher DD, Gallaher CM, Mahrt GJ, et al. A glucomannan and chitosan fiber supplement decreases plasma cholesterol and

- increases cholesterol excretion in overweight normocholesterolemic humans. *J Am Coll Nutr* 2002;21(5):428-33.
234. Gallaher CM, Munion J, Hesslink R, Jr., Wise J, Gallaher DD. Cholesterol reduction by glucomannan and chitosan is mediated by changes in cholesterol absorption and bile acid and fat excretion in rats. *J Nutr* 2000;130(11):2753-9.
 235. Chiang MT, Yao HT, Chen HC. Effect of dietary chitosans with different viscosity on plasma lipids and lipid peroxidation in rats fed on a diet enriched with cholesterol. *Biosci Biotechnol Biochem* 2000;64(5):965-71.
 236. Tai TS, Sheu WH, Lee WJ, Yao HT, Chiang MT. Effect of chitosan on plasma lipoprotein concentrations in type 2 diabetic subjects with hypercholesterolemia. *Diabetes Care* 2000;23(11):1703-4.
 237. Wuolijoki E, Hirvela T, Ylitalo P. Decrease in serum LDL cholesterol with microcrystalline chitosan. *Methods Find Exp Clin Pharmacol* 1999;21(5):357-61.
 238. Gueriolini R, Radu-Radulescu L, Boldrin M, Dallas J, Moore R. Comparative evaluation of fecal fat excretion induced by orlistat and chitosan. *Obes Res* 2001;9(6):364-7.
 239. Pittler MH, Abbot NC, Harkness EF, Ernst E. Randomized, double-blind trial of chitosan for body weight reduction. *Eur J Clin Nutr* 1999;53(5):379-81.
 240. Ho SC, Tai ES, Eng PH, Tan CE, Fok AC. In the absence of dietary surveillance, chitosan does not reduce plasma lipids or obesity in hypercholesterolaemic obese Asian subjects. *Singapore Med J* 2001;42(1):006-10.
 241. Penzak SR, Jann MW, Cold JA, Hon YY, Desai HD, Gurley BJ. Seville (sour) orange juice: synephrine content and cardiovascular effects in normotensive adults. *J Clin Pharmacol* 2001;41(10):1059-63.
 242. Moro CO, Basile G. Obesity and medicinal plants. *Fitoterapia* 2000;71 Suppl 1:S73-82.
 243. Cavallo E, Armellini F, Zamboni M, Vicentini R, Milani MP, Bosello O. Resting metabolic rate, body composition and thyroid hormones. Short term effects of very low calorie diet. *Horm Metab Res* 1990;22(12):632-5.
 244. Goglia F, Silvestri E, Lanni A. Thyroid hormones and mitochondria. *Biosci Rep* 2002;22(1):17-32.
 245. Wilson JH, Lamberts SW. The effect of triiodothyronine on weight loss and nitrogen balance of obese patients on a very-low-calorie liquid-formula diet. *Int J Obes* 1981;5(3):279-82.
 246. Rama Rao SV, Sunder GS, Reddy MR, Praharaj NK, Raju MV, Panda AK. Effect of supplementary choline on the performance of broiler breeders fed on different energy sources. *Br Poult Sci* 2001;42(3):362-7.
 247. Buchman AL, Awal M, Jenden D, Roch M, Kang SH. The effect of lecithin supplementation on plasma choline concentrations during a marathon. *J Am Coll Nutr* 2000;19(6):768-70.
 248. Buchman AL, Jenden D, Roch M. Plasma free, phospholipid-bound and urinary free choline all decrease during a marathon run and may be associated with impaired performance. *J Am Coll Nutr* 1999;18(6):598-601.
 249. Garcia Neto M, Pesti GM, Bakalli RI. Influence of dietary protein level on the broiler chicken's response to methionine and betaine supplements. *Poult Sci* 2000;79(10):1478-84.
 250. Overland M, Rorvik KA, Skrede A. Effect of trimethylamine oxide and betaine in swine diets on growth performance, carcass characteristics, nutrient digestibility, and sensory quality of pork. *J Anim Sci* 1999;77(8):2143-53.
 251. Schwab U, Torronen A, Toppinen L, et al. Betaine supplementation decreases plasma homocysteine concentrations but does not affect body weight, body composition, or resting energy expenditure in human subjects. *Am J Clin Nutr* 2002;76(5):961-7.
 252. Ammon HP, Muller AB. Forskolin: from an ayurvedic remedy to a modern agent. *Planta Med* 1985(6):473-7.
 253. Ammon HP, Muller AB. Effect of forskolin on islet cyclic AMP, insulin secretion, blood glucose and intravenous glucose tolerance in rats. *Naunyn Schmiedebergs Arch Pharmacol* 1984;326(4):364-7.
 254. de Souza NJ, Dohadwalla AN, Reden J. Forskolin: a labdane diterpenoid with antihypertensive, positive inotropic, platelet aggregation inhibitory, and adenylate cyclase activating properties. *Med Res Rev* 1983;3(2):201-19.
 255. Litosch I, Saito Y, Fain JN. Forskolin as an activator of cyclic AMP accumulation and secretion in blowfly salivary glands. *Biochem J* 1982;204(1):147-51.
 256. Litosch I, Hudson TH, Mills I, Li SY, Fain JN. Forskolin as an activator of cyclic AMP accumulation and lipolysis in rat adipocytes. *Mol Pharmacol* 1982;22(1):109-15.
 257. Seamon KB, Padgett W, Daly JW. Forskolin: unique diterpene activator of adenylate cyclase in membranes and in intact cells. *Proc Natl Acad Sci U S A* 1981;78(6):3363-7.
 258. Badmaev V, Majeed M, Conte AA, Parker JE. Diterpene forskolin (*Coleus forskohlii*, Benth.): A possible new compound for reduction of body weight by increasing lean body mass. Piscataway, NJ: Sabinsa Corporation; 2001.
 259. Kreider RB, Henderson S, Magu B, et al. Effects of *coleus forskohlii* supplementation on body composition and markers of health in sedentary overweight females. *FASEB J* 2002;LB59.
 260. Ebeling P, Koivisto VA. Physiological importance of dehydroepiandrosterone. *Lancet* 1994;343(8911):1479-81.
 261. Denti L, Pasolini G, Sanfelici L, et al. Effects of aging on dehydroepiandrosterone sulfate in relation to fasting insulin levels and body composition assessed by bioimpedance analysis. *Metabolism* 1997;46(7):826-32.
 262. De Pergola G, Zamboni M, Sciaraffia M, et al. Body fat accumulation is possibly responsible for lower dehydroepiandrosterone circulating levels in premenopausal obese women. *Int J Obes Relat Metab Disord* 1996;20(12):1105-10.
 263. Nestler JE, Barlascini CO, Clore JN, Blackard WG. Dehydroepiandrosterone reduces serum low density lipoprotein levels and body fat but does not alter insulin sensitivity in normal men. *J Clin Endocrinol Metab* 1988;66(1):57-61.
 264. Vogiatzi MG, Bueck MA, Vlachopapadopoulou E, el-Rashid R, New MI. Dehydroepiandrosterone in morbidly obese adolescents: effects on weight, body composition, lipids, and insulin resistance. *Metabolism* 1996;45(8):1011-5.
 265. Kalman DS, Colker CM, Swain MA, Torina GC, Shi Q. A randomized double-blind, placebo-controlled study of 3-acetyl-7-oxo-dehydroepiandrosterone in healthy overweight adults. *Curr Thera* 2000;61:435-42.
 266. MacDonald HB. Conjugated linoleic acid and disease prevention: a review of current knowledge. *J Am Coll Nutr* 2000;19(2 Suppl):111S-8S.
 267. Lowery LM, Appicelli PA, P.W.R. L. Conjugated linoleic acid enhances muscle size and strength gains in novice bodybuilders. *Med Sci Sports Exerc* 1998;30(5):S182.
 268. Riserus U, Berglund L, Vessby B. Conjugated linoleic acid (CLA) reduced abdominal adipose tissue in obese middle-aged men with signs of the metabolic syndrome: a randomised controlled trial. *Int J Obes Relat Metab Disord* 2001;25(8):1129-35.
 269. Riserus U, Arner P, Brismar K, Vessby B. Treatment with dietary trans10cis12 conjugated linoleic acid causes isomer-specific insulin resistance in obese men with the metabolic syndrome. *Diabetes Care* 2002;25(9):1516-21.

270. Riserus U, Basu S, Jovinge S, Fredrikson GN, Arnlov J, Vessby B. Supplementation with conjugated linoleic acid causes isomer-dependent oxidative stress and elevated C-reactive protein: a potential link to fatty acid-induced insulin resistance. *Circulation* 2002;106(15):1925-9.
271. Thom E, Wadstein J, Gudmundsen O. Conjugated linoleic acid reduces body fat in healthy exercising humans. *J Int Med Res* 2001;29(5):392-6.
272. Von Loeffelholz C, Von Loeffelholz BA, Von Loeffelholz B, Jahreis G. Influence of conjugated linoleic acid (CLA) supplementation on body composition and strength in bodybuilders. In: Symposium: Vitamine und Zusatzstoffe; 1999: Jena (Thhr.); 1999. p. 238-43.
273. Beuker F, Haak H, Schwietz H. CLA and body styling. In: Symposium: Vitamine und Zusatzstoffe; 1999: Jena (Thhr.); 1999. p. 229-37.
274. Medina EA, Horn WF, Keim NL, et al. Conjugated linoleic acid supplementation in humans: effects on circulating leptin concentrations and appetite. *Lipids* 2000;35(7):783-8.
275. Jena BS, Jayaprakasha GK, Singh RP, Sakariah KK. Chemistry and biochemistry of (-)-hydroxycitric acid from *Garcinia*. *J Agric Food Chem* 2002;50(1):10-22.
276. Ishihara K, Oyaizu S, Onuki K, Lim K, Fushiki T. Chronic (-)-hydroxycitrate administration spares carbohydrate utilization and promotes lipid oxidation during exercise in mice. *J Nutr* 2000;130(12):2990-5.
277. Kriketos AD, Thompson HR, Greene H, Hill JO. (-)-Hydroxycitric acid does not affect energy expenditure and substrate oxidation in adult males in a post-absorptive state. *Int J Obes Relat Metab Disord* 1999;23(8):867-73.
278. Heymsfield SB, Allison DB, Vasselli JR, Pirotbelli A, Greenfield D, Nunez C. *Garcinia cambogia* (hydroxycitric acid) as a potential antiobesity agent: a randomized controlled trial. *Jama* 1998;280(18):1596-600.
279. Mattes RD, Bormann L. Effects of (-)-hydroxycitric acid on appetitive variables. *Physiol Behav* 2000;71(1-2):87-94.
280. Brass EP. Supplemental carnitine and exercise. *Am J Clin Nutr* 2000;72(2 Suppl):618S-23S.
281. Villani RG, Gannon J, Self M, Rich PA. L-Carnitine supplementation combined with aerobic training does not promote weight loss in moderately obese women. *Int J Sport Nutr Exerc Metab* 2000;10(2):199-207.
282. Grases F, Melero G, Costa-Bauza A, Prieto R, March JG. Urolithiasis and phytotherapy. *Int Urol Nephrol* 1994;26(5):507-11.
283. Grases F, Llompert I, Conte A, Coll R, March JG. Glycosaminoglycans and oxalocalcic urolithiasis. *Nephron* 1994;68(4):449-53.
284. Dolan RL, Crosby EC, Leutkemeir MJ, Barton RG, Askew E, W. The effects of diuretics on resting metabolic rate and subsequent shifts in respiratory exchange ratios. *Med Sci Sports Exerc* 2001;33:S163.
285. Crosby EC, Dolan RL, Benson JE, Leutkemeir MJ, Barton RG, Askew EW. Herbal diuretic induced dehydration and resting metabolic rate. *Med Sci Sports Exerc* 2001;33:S163.
286. Derave W, Op't Eijnde B, Richter EA, Hespel P. Combined creatine and protein supplementation improves glucose tolerance and muscle glycogen accumulation in humans. Abstracts of 6th International Conference on Guanidino Compounds in Biology and Medicine 2001.
287. Nelson AG, Arnall DA, Kokkonen J, Day, Evans J. Muscle glycogen supercompensation is enhanced by prior creatine supplementation. *Med Sci Sports Exerc* 2001;33(7):1096-100.
288. Op't Eijnde B, Urso B, Richter EA, Greenhaff PL, Hespel P. Effect of oral creatine supplementation on human muscle GLUT4 protein content after immobilization. *Diabetes* 2001;50(1):18-23.
289. Green AL, Hultman E, Macdonald IA, Sewell DA, Greenhaff PL. Carbohydrate ingestion augments skeletal muscle creatine accumulation during creatine supplementation in humans. *Am J Physiol* 1996;271(5 Pt 1):E821-6.
290. Nelson AG, Day R, Glickman-Weiss EL, Hegsted M, Kokkonen J, Sampson B. Creatine supplementation alters the response to a graded cycle ergometer test. *Eur J Appl Physiol* 2000;83(1):89-94.
291. Nelson AG, Day R, Glickman-Weiss EL, Hegsted M, Sampson B. Creatine supplementation raises anaerobic threshold. *FASEB Journal* 1997;11:A589.
292. Cade R, Conte M, Zauner C, et al. Effects of phosphate loading on 2,3 diphosphoglycerate and maximal oxygen uptake. *Med Sci Sports Exerc* 1984;16:263-8.
293. Stewart I, McNaughton L, Davies P, Tristram S. Phosphate loading and the effects of VO₂max in trained cyclists. *Res Quart* 1990;61:80-4.
294. McNaughton L, Backx K, Palmer G, Strange N. Effects of chronic bicarbonate ingestion on the performance of high-intensity work. *Eur J Appl Physiol Occup Physiol* 1999;80(4):333-6.
295. Applegate E. Effective nutritional ergogenic aids. *Int J Sport Nutr* 1999;9(2):229-39.
296. Kronfeld DS, Ferrante PL, Grandjean D. Optimal nutrition for athletic performance, with emphasis on fat adaptation in dogs and horses. *J Nutr* 1994;124(12 Suppl):2745S-53S.
297. Kraemer WJ, Gordon SE, Lynch JM, Pop ME, Clark KL. Effects of multibuffer supplementation on acid-base balance and 2,3- diphosphoglycerate following repetitive anaerobic exercise. *Int J Sport Nutr* 1995;5(4):300-14.
298. Matson LG, Tran ZV. Effects of sodium bicarbonate ingestion on anaerobic performance: a meta-analytic review. *Int J Sport Nutr* 1993;3(1):2-28.
299. Graham TE. Caffeine, coffee and ephedrine: impact on exercise performance and metabolism. *Can J Appl Physiol* 2001;26(Suppl):S103-19.
300. Tarnopolsky MA, Atkinson SA, MacDougall JD, Sale DG, Sutton JR. Physiological responses to caffeine during endurance running in habitual caffeine users. *Med Sci Sports Exerc* 1989;21(4):418-24.
301. Armstrong LE. Caffeine, body fluid-electrolyte balance, and exercise performance. *Int J Sport Nutr Exerc Metab* 2002;12(2):189-206.
302. Graham TE. Caffeine and exercise: metabolism, endurance and performance. *Sports Med* 2001;31(11):785-807.
303. Falk B, Burstein R, Rosenblum J, Shapiro Y, Zylber-Katz E, Bashan N. Effects of caffeine ingestion on body fluid balance and thermoregulation during exercise. *Can J Physiol Pharmacol* 1990;68(7):889-92.
304. Tarnopolsky MA. Potential use of creatine monohydrate in muscular dystrophy and neurometabolic disorders. Abstracts of 6th International Conference on Guanidino Compounds in Biology and Medicine 2001.
305. Wagner DR. Hyperhydrating with glycerol: implications for athletic performance. *J Am Diet Assoc* 1999;99(2):207-12.
306. Inder WJ, Swanney MP, Donald RA, Prickett TC, Hellemans J. The effect of glycerol and desmopressin on exercise performance and hydration in triathletes. *Med Sci Sports Exerc* 1998;30(8):1263-9.

307. Montner P, Stark DM, Riedesel ML, et al. Pre-exercise glycerol hydration improves cycling endurance time. *Int J Sports Med* 1996;17(1):27-33.
308. Boulay MR, Song TM, Serresse O, Theriault G, Simoneau JA, Bouchard C. Changes in plasma electrolytes and muscle substrates during short-term maximal exercise in humans. *Can J Appl Physiol* 1995;20(1):89-101.
309. Tikuisis P, Ducharme MB, Moroz D, Jacobs I. Physiological responses of exercised-fatigued individuals exposed to wet-cold conditions. *J Appl Physiol* 1999;86(4):1319-28.
310. Jimenez C, Melin B, Koulmann N, Allevard AM, Launay JC, Savourey G. Plasma volume changes during and after acute variations of body hydration level in humans. *Eur J Appl Physiol Occup Physiol* 1999;80(1):1-8.
311. Magal M, Webster MJ, Sistrunk LE, Whitehead MT, Evans RK, Boyd JC. Comparison of glycerol and water hydration regimens on tennis-related performance. *Med Sci Sports Exerc* 2003;35(1):150-6.
312. Bell DG, McLellan TM, Sabiston CM. Effect of ingesting caffeine and ephedrine on 10-km run performance. *Med Sci Sports Exerc* 2002;34(2):344-9.
313. Bell DG, Jacobs I, Ellerington K. Effect of caffeine and ephedrine ingestion on anaerobic exercise performance. *Med Sci Sports Exerc* 2001;33(8):1399-403.
314. Bell DG, Jacobs I, McLellan TM, Zamecnik J. Reducing the dose of combined caffeine and ephedrine preserves the ergogenic effect. *Aviat Space Environ Med* 2000;71(4):415-9.
315. Bell DG, Jacobs I. Combined caffeine and ephedrine ingestion improves run times of Canadian Forces Warrior Test. *Aviat Space Environ Med* 1999;70(4):325-9.
316. Bell DG, Jacobs I, Zamecnik J. Effects of caffeine, ephedrine and their combination on time to exhaustion during high-intensity exercise. *Eur J Appl Physiol Occup Physiol* 1998;77(5):427-33.
317. Jeukendrup AE, Saris WH, Wagenmakers AJ. Fat metabolism during exercise: a review--part III: effects of nutritional interventions. *Int J Sports Med* 1998;19(6):371-9.
318. Goedecke JH, Elmer-English R, Dennis SC, Schloss I, Noakes TD, Lambert EV. Effects of medium-chain triacylglycerol ingested with carbohydrate on metabolism and exercise performance. *Int J Sport Nutr* 1999;9(1):35-47.
319. Calabrese C, Myer S, Munson S, Turet P, Birdsall TC. A cross-over study of the effect of a single oral feeding of medium chain triglyceride oil vs. canola oil on post-ingestion plasma triglyceride levels in healthy men. *Altern Med Rev* 1999;4(1):23-8.
320. Angus DJ, Hargreaves M, Dancy J, Febbraio MA. Effect of carbohydrate or carbohydrate plus medium-chain triglyceride ingestion on cycling time trial performance. *J Appl Physiol* 2000;88(1):113-9.
321. Jeukendrup AE, Thielen JJ, Wagenmakers AJ, Brouns F, Saris WH. Effect of medium-chain triacylglycerol and carbohydrate ingestion during exercise on substrate utilization and subsequent cycling performance. *Am J Clin Nutr* 1998;67(3):397-404.
322. Van Zyl CG, Lambert EV, Hawley JA, Noakes TD, Dennis SC. Effects of medium-chain triglyceride ingestion on fuel metabolism and cycling performance. *J Appl Physiol* 1996;80(6):2217-25.
323. Tullson PC, Terjung RL. Adenine nucleotide synthesis in exercising and endurance-trained skeletal muscle. *Am J Physiol* 1991;261(2 Pt 1):C342-7.
324. Gross M, Kormann B, Zollner N. Ribose administration during exercise: effects on substrates and products of energy metabolism in healthy subjects and a patient with myoadenylate deaminase deficiency. *Klin Wochenschr* 1991;69(4):151-5.
325. Wagner DR, Gresser U, Kamilli I, Gross M, Zollner N. Effects of oral ribose on muscle metabolism during bicycle ergometer in patients with AMP-deaminase-deficiency. *Adv Exp Med Biol* 1991:383-5.
326. Pliml W, von Arnim T, Stablein A, Hofmann H, Zimmer HG, Erdmann E. Effects of ribose on exercise-induced ischaemia in stable coronary artery disease. *Lancet* 1992;340(8818):507-10.
327. Pauly DF, Pepine CJ. D-Ribose as a supplement for cardiac energy metabolism. *J Cardiovasc Pharmacol Ther* 2000;5(4):249-58.
328. Op 't Eijnde B, Van Leemputte M, Brouns F, et al. No effects of oral ribose supplementation on repeated maximal exercise and de novo ATP resynthesis. *J Appl Physiol* 2001;91(5):2275-81.
329. Berardi JM, Ziegenfuss TN. Effects of ribose supplementation on repeated sprint performance in men. *J Strength Cond Res* 2003;17(1):47-52.
330. Kreider RB, Melton C, Greenwood M, et al. Effects of oral d-ribose supplementation on anaerobic capacity and selected metabolic markers in healthy males. *Int J Sport Nutr Exerc Metab* 2003;13(1):87-96.
331. Hargreaves M, McKenna MJ, Jenkins DG, et al. Muscle metabolites and performance during high-intensity, intermittent exercise. *J Appl Physiol* 1998;84(5):1687-91.
332. Starling RD, Trappe TA, Short KR, et al. Effect of inosine supplementation on aerobic and anaerobic cycling performance. *Med Sci Sports Exerc* 1996;28(9):1193-8.
333. Williams MH, Kreider RB, Hunter DW, et al. Effect of inosine supplementation on 3-mile treadmill run performance and VO₂ peak. *Med Sci Sports Exerc* 1990;22(4):517-22.
334. McNaughton L, Dalton B, Tarr J. Inosine supplementation has no effect on aerobic or anaerobic cycling performance. *Int J Sport Nutr* 1999;9(4):333-44.
335. Braham R, Dawson B, Goodman C. The effect of glucosamine supplementation on people experiencing regular knee pain. *Br J Sports Med* 2003;37(1):45-9; discussion 9.
336. Vad V, Hong HM, Zazzali M, Agi N, Basrai D. Exercise recommendations in athletes with early osteoarthritis of the knee. *Sports Med* 2002;32(11):729-39.
337. Nieman DC. Exercise immunology: nutritional countermeasures. *Can J Appl Physiol* 2001;26(Suppl):S45-55.
338. Gleeson M, Lancaster GI, Bishop NC. Nutritional strategies to minimise exercise-induced immunosuppression in athletes. *Can J Appl Physiol* 2001;26(Suppl):S23-35.
339. Gleeson M, Bishop NC. Elite athlete immunology: importance of nutrition. *Int J Sports Med* 2000;21 Suppl 1:S44-50.
340. Lowery L, Berardi JM, Ziegenfuss T. Antioxidants. In: Antonio J, Stout J, eds. *Sports Supplements*. Baltimore, MD: Lippincott, Williams & Wilkins; 2001:260-78.
341. Gomez AL, Volek JS, Ratamess NA, et al. Creatine supplementation enhances body composition during short-term resistance training overreaching. *Journal of Strength and Conditioning Research* 2000;14(3).
342. French DN, Volek JS, Ratamess NA, et al. The effects of creatine supplementation on resting serum hormonal concentrations during short-term resistance training overreaching. *Med Sci Sports & Exerc* 2001;33(5):S203.

343. Volek JS, Kraemer WJ, Rubin MR, Gomez AL, Ratamess NA, Gaynor P. L-Carnitine L-tartrate supplementation favorably affects markers of recovery from exercise stress. *Am J Physiol Endocrinol Metab* 2002;282(2):E474-82.
344. Mero A. Leucine supplementation and intensive training. *Sports Med* 1999;27(6):347-58.
345. Harris RC, Soderlund K, Hultman E. Elevation of creatine in resting and exercised muscle of normal subjects by creatine supplementation. *Clin Sci (Colch)* 1992;83(3):367-74.
346. Williams MH. Vitamin supplementation and athletic performance. *Int J Vitam Nutr Res Suppl* 1989;30(163-91).
347. Reid IR. Therapy of osteoporosis: calcium, vitamin D, and exercise. *Am J Med Sci* 1996;312(6):278-86.
348. Goldfarb AH. Antioxidants: role of supplementation to prevent exercise-induced oxidative stress. *Med Sci Sports Exerc* 1993;25(2):232-6.
349. Goldfarb AH. Nutritional antioxidants as therapeutic and preventive modalities in exercise-induced muscle damage. *Can J Appl Physiol* 1999;24(3):249-66.
350. Appell HJ, Duarte JA, Soares JM. Supplementation of vitamin E may attenuate skeletal muscle immobilization atrophy. *Int J Sports Med* 1997;18(3):157-60.
351. Tiidus PM, Houston ME. Vitamin E status and response to exercise training. *Sports Med* 1995;20(1):12-23.
352. Craciun AM, Wolf J, Knapen MH, Brouns F, Vermeer C. Improved bone metabolism in female elite athletes after vitamin K supplementation. *Int J Sports Med* 1998;19(7):479-84.
353. Fogelholm M, Ruokonen I, Laakso JT, Vuorimaa T, Himberg JJ. Lack of association between indices of vitamin B1, B2, and B6 status and exercise-induced blood lactate in young adults. *Int J Sport Nutr* 1993;3(2):165-76.
354. Garg R, Malinow M, Pettinger M, Upson B, Hunninghake D. Niacin treatment increases plasma homocyst(e)ine levels. *Am Heart J* 1999;138(6 Pt 1):1082-7.
355. Alaswad K, O'Keefe JH, Jr., Moe RM. Combination drug therapy for dyslipidemia. *Curr Atheroscler Rep* 1999;1(1):44-9.
356. Murray R, Bartoli WP, Eddy DE, Horn MK. Physiological and performance responses to nicotinic-acid ingestion during exercise. *Med Sci Sports Exerc* 1995;27(7):1057-62.
357. Bonke D. Influence of vitamin B1, B6, and B12 on the control of fine motoric movements. *Bibl Nutr Dieta* 1986(38):104-9.
358. Bonke D, Nickel B. Improvement of fine motoric movement control by elevated dosages of vitamin B1, B6, and B12 in target shooting. *Int J Vitam Nutr Res Suppl* 1989;30:198-204.
359. Van Dyke DC, Stumbo PJ, Mary JB, Niebyl JR. Folic acid and prevention of birth defects. *Dev Med Child Neurol* 2002;44(6):426-9.
360. Mattson MP, Kruman, II, Duan W. Folic acid and homocysteine in age-related disease. *Ageing Res Rev* 2002;1(1):95-111.
361. Weston PM, King RF, Goode AW, Williams NS. Diet-induced thermogenesis in patients with gastrointestinal cancer cachexia. *Clin Sci (Lond)* 1989;77(2):133-8.
362. Webster MJ. Physiological and performance responses to supplementation with thiamin and pantothenic acid derivatives. *Eur J Appl Physiol Occup Physiol* 1998;77(6):486-91.
363. van der Beek EJ, Lowik MR, Hulshof KF, Kistemaker C. Combinations of low thiamin, riboflavin, vitamin B6 and vitamin C intake among Dutch adults. (Dutch Nutrition Surveillance System). *J Am Coll Nutr* 1994;13(4):383-91.
364. van der Beek EJ. Vitamin supplementation and physical exercise performance. *J Sports Sci* 1991;9 Spec No:77-90.
365. Pedersen BK, Bruunsgaard H, Jensen M, Krzykowski K, Ostrowski K. Exercise and immune function: effect of ageing and nutrition. *Proc Nutr Soc* 1999;58(3):733-42.
366. Petersen EW, Ostrowski K, Ibfelt T, et al. Effect of vitamin supplementation on cytokine response and on muscle damage after strenuous exercise. *Am J Physiol Cell Physiol* 2001;280(6):C1570-5.
367. Grados F, Brazier M, Kamel S, et al. Effects on bone mineral density of calcium and vitamin D supplementation in elderly women with vitamin D deficiency. *Joint Bone Spine* 2003;70(3):203-8.
368. Campbell WW, Joseph LJ, Anderson RA, Davey SL, Hinton J, Evans WJ. Effects of resistive training and chromium picolinate on body composition and skeletal muscle size in older women. *Int J Sport Nutr Exerc Metab* 2002;12(2):125-35.
369. Brutsaert TD, Hernandez-Cordero S, Rivera J, Viola T, Hughes G, Haas JD. Iron supplementation improves progressive fatigue resistance during dynamic knee extensor exercise in iron-depleted, nonanemic women. *Am J Clin Nutr* 2003;77(2):441-8.
370. Bohl CH, Volpe SL. Magnesium and exercise. *Crit Rev Food Sci Nutr* 2002;42(6):533-63.
371. Lukaski HC. Magnesium, zinc, and chromium nutrition and athletic performance. *Can J Appl Physiol* 2001;26 Suppl:S13-22.
372. Morton DP, Callister R. Characteristics and etiology of exercise-related transient abdominal pain. *Med Sci Sports Exerc* 2000;32(2):432-8.
373. Noakes TD. Fluid and electrolyte disturbances in heat illness. *Int J Sports Med* 1998;19 Suppl 2:S146-9.
374. Sawka MN, Montain SJ. Fluid and electrolyte supplementation for exercise heat stress. *Am J Clin Nutr* 2000;72(2 Suppl):564S-72S.
375. Margaritis I, Tessier F, Prou E, Marconnet P, Marini JF. Effects of endurance training on skeletal muscle oxidative capacities with and without selenium supplementation. *J Trace Elem Med Biol* 1997;11(1):37-43.
376. Tessier F, Margaritis I, Richard MJ, Moynot C, Marconnet P. Selenium and training effects on the glutathione system and aerobic performance. *Med Sci Sports Exerc* 1995;27(3):390-6.
377. McCutcheon LJ, Geor RJ. Sweating. Fluid and ion losses and replacement. *Vet Clin North Am Equine Pract* 1998;14(1):75-95.
378. Gibson RS, Heath AL, Ferguson EL. Risk of suboptimal iron and zinc nutriture among adolescent girls in Australia and New Zealand: causes, consequences, and solutions. *Asia Pac J Clin Nutr* 2002;11 Suppl 3:S543-52.
379. Singh A, Failla ML, Deuster PA. Exercise-induced changes in immune function: effects of zinc supplementation. *J Appl Physiol* 1994;76(6):2298-303.

Table 1. Proposed Nutritional Ergogenic Aids – Vitamins

Nutrient	RDA	Proposed Ergogenic Value	Summary of Research Findings
Vitamin A	Males 900 mcg/d Females 700 mcg/d	Constituent of rhodopsin (visual pigment) and is involved in night vision. Some suggest that vitamin A supplementation may improve sport vision.	No studies have shown that vitamin A supplementation improves exercise performance ³⁴⁶ .
Vitamin D	5 mcg/d (age <51)	Promotes bone growth and mineralization. Enhances calcium absorption. Supplementation with calcium may help prevent bone loss in osteoporotic populations.	Co-supplementation with calcium may help prevent bone loss in athletes susceptible to osteoporosis ³⁴⁷ . However, vitamin D supplementation does not enhance exercise performance ³⁴⁶ .
Vitamin E	15 mg/d	As an antioxidant, it has been shown to help prevent the formation of free radicals during intense exercise and prevent the destruction of red blood cells, improving or maintaining oxygen delivery to the muscles during exercise. Some evidence suggests that it may reduce risk to heart disease or decrease incidence of recurring heart attack.	Numerous studies show that vitamin E supplementation can decrease exercise-induced oxidative stress ³⁴⁸⁻³⁵⁰ . However, most studies show no effects on performance at sea level. At high altitudes, vitamin E may improve exercise performance ³⁵¹ . Additional research is necessary to determine whether long-term supplementation may help athletes better tolerate training.
Vitamin K	Males 120 mcg/d Females 90 mcg/d	Important in blood clotting. There is also some evidence that it may affect bone metabolism in postmenopausal women.	Vitamin K supplementation (10 mg/d) in elite female athletes has been reported to increase calcium-binding capacity of osteocalcin and promoted a 15-20% increase in bone formation markers and a 20-25% decrease in bone resorption markers suggesting an improved balance between bone formation and resorption ³⁵² .
Thiamin (B ₁)	Males 1.2 mg/d Females 1.1 mg/d	Coenzyme (thiamin pyrophosphate) in the removal of CO ₂ from decarboxylic reactions from pyruvate to acetyl CoA and in TCA cycle. Supplementation is theorized to improve anaerobic threshold and CO ₂ transport. Deficiencies may decrease efficiency of energy systems.	Dietary availability of thiamin does not appear to affect exercise capacity when athletes have a normal intake ³⁵³ .
Riboflavin (B ₂)	Males 1.3 mg/d Females 1.7 mg/d	Constituent of flavin nucleotide coenzymes involved in energy metabolism. Theorized to enhance energy availability during oxidative metabolism.	Dietary availability of riboflavin does not appear to affect exercise capacity when athletes have a normal intake ³⁵³ .
Niacin (B ₃)	Males 16 mg/d Females 14 mg/d	Constituent of coenzymes involved in energy metabolism. Theorized to blunt increases in fatty acids during exercise, reduce cholesterol, enhance thermoregulation, and improve energy availability during oxidative metabolism.	Studies indicate that niacin supplementation (100-500 mg/d) can help decrease blood lipid levels and increase homocysteine levels in hypercholesteremic patients ^{354, 355} . However, niacin supplementation (280 mg) during exercise has been reported to decrease exercise capacity by blunting the mobilization of fatty acids ³⁵⁶ .
Pyridoxine (B ₆)	1.3 mg/d (age <51)	Has been marketed as a supplement that will improve muscle mass, strength, and aerobic power in the lactic acid and oxygen systems. It also may have a calming effect that has been linked to an improved mental strength.	In well-nourished athletes, pyridoxine failed to improve aerobic capacity, or lactic acid accumulation ³⁵³ . However, when combined with vitamins B ₁ and B ₁₂ , it may increase serotonin levels and improve fine motor skills that may be necessary in sports like pistol shooting and archery ^{357, 358} .

Cyano-cobalamin (B ₁₂)	2.4 mcg/d	A coenzyme involved in the production of DNA and serotonin. DNA is important in protein and red blood cell synthesis. Theoretically, it would increase muscle mass, the oxygen-carrying capacity of blood, and decrease anxiety.	In well-nourished athletes, no ergogenic effect has been reported. However, when combined with vitamins B ₁ and B ₆ , cyanocobalamin has been shown to improve performance in pistol shooting ³⁵⁸ . This may be due to increased levels of serotonin, a neurotransmitter in the brain, which may reduce anxiety.
Folic acid (folate)	400 mcg/d	Functions as a coenzyme in the formation of DNA and red blood cells. An increase in red blood cells could improve oxygen delivery to the muscles during exercise. Believed to be important to help prevent birth defects and may help decrease homocysteine levels.	Studies suggest that increasing dietary availability of folic acid during pregnancy can lower the incidence of birth defects ³⁵⁹ . Additionally, it may decrease homocysteine levels (a risk factor for heart disease) ³⁶⁰ . In well-nourished and folate deficient-athletes, folic acid did not improve exercise performance ³⁶¹ .
Pantothenic acid	5 mg/d	Acts as a coenzyme for acetyl coenzyme A (acetyl CoA). This may benefit aerobic or oxygen energy systems.	Research has reported no improvements in aerobic performance with acetyl CoA supplementation. However, one study reported a decrease in lactic acid accumulation, without an improvement in performance ³⁶² .
Beta carotene	None	Serves as an antioxidant. Theorized to help minimize exercise-induced lipid peroxidation and muscle damage.	Research indicates that beta carotene supplementation with or without other antioxidants can help decrease exercise-induced peroxidation. Over time, this may help athletes tolerate training. However, it is unclear whether antioxidant supplementation affects exercise performance ³⁴⁹ .
Vitamin C	Males 90 mg/d Females 75 mg/d	Used in a number of different metabolic processes in the body. It is involved in the synthesis of epinephrine, iron absorption, and is an antioxidant. Theoretically, it could benefit exercise performance by improving metabolism during exercise. There is also evidence that vitamin C may enhance immunity.	In well-nourished athletes, vitamin C supplementation does not appear to improve physical performance ^{363, 364} . However, there is some evidence that vitamin C supplementation (e.g., 500 mg/d) following intense exercise may decrease the incidence of upper respiratory tract infections ^{337, 365, 366} .

Recommended Dietary Allowances (RDA) based on the 1989 Food & Nutrition Board, National Academy of Sciences-National Research Council recommendations. Updated in 2001.

Table 2. Proposed Nutritional Ergogenic Aids – Minerals

Nutrient	RDA	Proposed Ergogenic Value	Summary of Research Findings
Boron	None	Boron has been marketed to athletes as a dietary supplement that may promote muscle growth during resistance training. The rationale was primarily based on an initial report that boron supplementation (3 mg/d) significantly increased β -estradiol and testosterone levels in postmenopausal women consuming a diet low in boron.	Studies which have investigated the effects of 7 wk of boron supplementation (2.5 mg/d) during resistance training on testosterone levels, body composition, and strength have reported no ergogenic value ^{126, 127} . There is no evidence at this time that boron supplementation during resistance-training promotes muscle growth.
Calcium	1000 mg/d (ages 19-50)	Involved in bone and tooth formation, blood clotting, and nerve transmission. Stimulates fat metabolism. Diet should contain sufficient amounts, especially in growing children/adole ¹²⁹ scents, female athletes, and postmenopausal women. Vitamin D needed to assist absorption.	Calcium supplementation may be beneficial in populations susceptible to osteoporosis ³⁶⁷ . Additionally, calcium supplementation has been shown to promote fat metabolism and help manage body composition ^{214, 215} . Calcium supplementation provides no ergogenic effect on exercise performance.
Chromium	Males 35 mcg/d Females 25 mcg/d (ages 19-50)	Chromium, commonly sold as chromium picolinate, has been marketed with claims that the supplement will increase lean body mass and decrease body fat levels.	Animal research indicates that chromium supplementation increases lean body mass and reduces body fat. Early research on humans reported similar results ¹²⁹ , however, more recent well-controlled studies reported that chromium supplementation (200 to 800 mcg/d) does not improve lean body mass or reduce body fat ^{134, 368} .
Iron	Males 8 mg/d Females 18 mg/d (age 19-50)	Iron supplements are used to increase aerobic performance in sports that use the oxygen system. Iron is a component of hemoglobin in the red blood cell, which is a carrier of oxygen.	Most research shows that iron supplements do not appear to improve aerobic performance unless the athlete is iron-depleted and/or has anemia ³⁶⁹ .
Magnesium	Males 420 Females 320	Activates enzymes involved in protein synthesis. Involved in ATP reactions. Serum levels decrease with exercise. Some suggest that magnesium supplementation may improve energy metabolism/ATP availability.	Most well-controlled research indicates that magnesium supplementation (500 mg/d) does not affect exercise performance in athletes unless there is a deficiency ^{370, 371} .
Phosphorus (phosphate salts)	700 mg/d	Phosphate has been studied for its ability to improve all three energy systems, primarily the oxygen system or aerobic capacity.	Recent well-controlled research studies reported that sodium phosphate supplementation (4 g/d for 3 d) improved the oxygen energy system in endurance tasks ^{219, 220, 292} . There appears to be little ergogenic value of other forms of phosphate (i.e., calcium phosphate, potassium phosphate). More research is needed to determine the mechanism for improvement.
Potassium	2000 mg/d*	An electrolyte that helps regulate fluid balance, nerve transmission, and acid-base balance. Some suggest excessive increases or decreases in potassium may predispose athletes to cramping.	Although potassium loss during intense exercise in the heat has been anecdotally associated with muscle cramping, the etiology of cramping is unknown ^{372, 373} . It is unclear whether potassium supplementation in athletes decreases the incidence of muscle cramping ³⁷⁴ . No ergogenic effects reported.

Selenium	55 mcg/d	Marketed as a supplement to increase aerobic exercise performance. Working closely with vitamin E and glutathione peroxidase (an antioxidant), selenium may destroy destructive free radical production of lipids during aerobic exercise.	Although selenium may reduce lipid peroxidation during aerobic exercise, improvements in aerobic capacity have not been demonstrated ^{375, 376} .
Sodium	500 mg/d*		During the first several days of intense training in the heat, a greater amount of sodium is lost in sweat. Additionally, prolonged ultraendurance exercise may decrease sodium levels leading to hyponatremia. Increasing salt availability during heavy training in the heat has been shown to help maintain fluid balance and prevent hyponatremia ^{374, 377} .
Vanadyl sulfate (vanadium)	None	Vanadium may be involved in reactions in the body that produce insulin-like effects on protein and glucose metabolism. Due to the anabolic nature of insulin, this has brought attention to vanadium as a supplement to increase muscle mass, enhance strength and power.	Limited research has shown that type 2 diabetics may improve their glucose control; however, there is no proof that vanadyl sulfate has any effect on muscle mass, strength, or power ^{189, 190} .
Zinc	Males 11 mg/d Females 8 mg/d	Constituent of enzymes involved in digestion. Associated with immunity. Theorized to reduce incidence of upper respiratory tract infections in athletes involved in heavy training.	Studies indicate that zinc supplementation (25 mg/d) during training minimized exercise-induced changes in immune function ^{42, 339, 378, 379} .

Recommended Dietary Allowances (RDA) based on the 2002 Food & Nutrition Board, National Academy of Sciences-National Research Council recommendations.

* Estimated minimum requirement

Table 3. Categorization of the Ergogenic Value of Performance Enhancement, Muscle Building, and Weight Loss Supplements

Category	Muscle Building Supplements	Weight Loss Supplements	Performance Enhancement
I. Apparently effective and generally safe	<ul style="list-style-type: none"> • Weight gain powders • Creatine • HMB (untrained individuals initiating training) 	<ul style="list-style-type: none"> • Low-calorie foods, MRPs and RTDs that help individuals maintain a hypocaloric diet • Ephedra, caffeine, and salicin-containing thermogenic supplements taken at recommended doses in appropriate populations (now banned by FDA) 	<ul style="list-style-type: none"> • Water and sports drinks • Carbohydrate • Creatine • Sodium phosphate • Sodium bicarbonate • Caffeine
II. Possibly effective	<ul style="list-style-type: none"> • Postexercise carbohydrate and protein • BCAA • Essential amino acids (EAA) • Glutamine • Protein • HMB (trained subjects) 	<ul style="list-style-type: none"> • High-fiber diets • Calcium • Phosphate • Green tea extract • Pyruvate/DHAP (at high doses) 	<ul style="list-style-type: none"> • Postexercise CHO/PRO • Glutamine • EAA • BCAA • HMB (trained subjects) • Glycerol • Low doses of ephedrine/caffeine (now banned by FDA)
III. Too early to tell	<ul style="list-style-type: none"> • α-Ketoglutarate • α-Ketoisocaproate (KIC) • Ecdysterones • Growth hormone releasing peptides (GHRP) and secretogues • HMB (trained athletes) • Isoflavones • Sulfo-polysaccharides (myostatin inhibitors) • Zinc/magnesium aspartate (ZMA) 	<ul style="list-style-type: none"> • Appetite suppressants and fat blockers (Gymnema sylvestre, chitosan) • Thermogenics (synephrine, thyroid stimulators, cayenne pepper, black pepper, ginger root) • Lipolytic nutrients (phosphatidyl choline, betaine, Coleus forskohlii, 7-keto DHEA) • Psychotropic Nutrients/Herbs 	<ul style="list-style-type: none"> • Medium chain triglycerides • Ribose
IV. Apparently not effective and/or dangerous	<ul style="list-style-type: none"> • Boron • Chromium • Conjugated linoleic acids (CLA) • Gamma oryzanol (ferulic acid) • Prohormones • Tribulus terrestris • Vanadyl sulfate (vanadium) • Yohimbe (Yohimbine) 	<ul style="list-style-type: none"> • Chromium (nondiabetics) • CLA • HCA • L-Carnitine • Pyruvate (at low doses) • Herbal diuretics • High doses of ephedrine/caffeine 	<ul style="list-style-type: none"> • Inosine • High doses of ephedrine/caffeine